

Systematic review: the evidence base for long-term management of coeliac disease

M. L. HAINES, R. P. ANDERSON & P. R. GIBSON

Department of Gastroenterology,
Monash University Department of
Medicine, Box Hill Hospital, Box Hill,
and Walter and Eliza Hall Institute,
Parkville, Vic., Australia

Correspondence to:

Prof. P. R. Gibson, Department of
Medicine, Box Hill Hospital, Box Hill,
Vic. 3128, Australia.
E-mail: peter.gibson@med.monash.
edu.au

Publication data

Submitted 1 June 2008
First decision 11 July 2008
Resubmitted 26 July 2008
Accepted 27 July 2008
Epub Accepted Article 30 July 2008

SUMMARY

Background

While gluten-free diet is an effective treatment for coeliac disease, the need for and goals of long-term management of patients are poorly defined.

Aim

To review systematically the complications and associations of coeliac disease, to identify potential risk factors, to define ways of assessing risk factors and to provide a strategy for management.

Methods

Review of medical literature from 1975.

Results

There is an increasing list of potential complications and/or conditions associated with coeliac disease, in particular, autoimmune disease, malignancy and bone disease. Risk factors that may predict or influence long-term outcomes include genetic susceptibility, environmental factors predominantly gluten ingestion, persistent small intestinal inflammation/injury and nutritional deficiencies. Genotyping of patients is yet to have an established clinical role in long-term management. Assessment of adherence to the gluten-free diet largely relies upon skilled dietary history, but the ultimate test is duodenal histopathology, which is the only currently established means of assessing healing. Symptoms, serology or other non-invasive means are poor predictors of healing and the likelihood of complications.

Conclusion

Evidence (albeit limited) that adherence to a gluten-free diet and mucosal healing prevent and/or ameliorate complications indicates that a planned long-term strategy for follow-up is essential.

Aliment Pharmacol Ther 28, 1042–1066

INTRODUCTION

The introduction of serological tests with good specificity for coeliac disease (CD) has led to the realization that this condition is common. In fact, the rate of diagnosis has markedly increased in recent years.^{1, 2} Treatment with a gluten-free diet (GFD) is successful in improving quality of life and in healing the small intestine in a majority of patients and recent detailed guidelines on the diet relevant to Australia and New Zealand have been published.³ However, it is becoming increasingly recognized that patients with CD are at increased risk of developing long-term complications and other autoimmune conditions. Can these be prevented and outcomes improved?

It is commonly argued by experienced clinicians that once the diagnosis is made, GFD instituted and symptoms improved, a majority of patients do not require repeat duodenal biopsy and medical follow-up is needed only where the symptoms attributed to the CD recur. While such an approach may be adequate for majority of patients, it may disadvantage the minority. On the other hand, those who claim that intense life-long follow-up with repeat duodenal biopsies, blood tests and other investigations may be accused of over-servicing a patient group and inducing a culture of the 'worried well'. To decide rationally where the ideal viewpoint rests, several issues require addressing. These include definition of the complications and consequences of CD; whether the risk of these relates to strict adherence to the GFD and/or healing of the small intestine or to neither; the best methods for assessing adherence to GFD and healing of the small intestine and the practical issue of the optimal way patients should be followed in routine clinical practice.

COMPLICATIONS AND CONSEQUENCES OF CD

There is a large body of literature describing conditions and complications associated with CD. These, in addition to the strength of the associations, are shown in Table 1.

RISK FACTORS FOR COMPLICATIONS

If risk factors for outcomes can be identified, then patients could be more rationally stratified in terms of the monitoring schedule applied and the use of preventive interventions. While there is a paucity of

definitive information, considerable data do exist and these are summarized in Table 2. Risk factors have been categorized into four potential areas: genetic, gluten exposure, intestinal inflammation and nutritional deficiency.

Genetic

Susceptibility to CD is determined to a significant extent by genetic factors. Approximately 88% of CD patients carry the HLA DQ2 heterodimer encoded by alleles DQA1*05 and DQB1*02.^{4, 5} The remainder possess either HLA DQA1*05 or HLA DQB1*02, or express HLA DQ8 encoded by alleles DQA1*03 and DQB1*0302.^{5, 6} HLA DQ2 and DQ8 are within haplotypes that are also associated with other autoimmune conditions, such as type-1 diabetes mellitus,⁷⁻⁹ autoimmune thyroid disease,¹⁰⁻¹² primary biliary cirrhosis,¹³ Addison's disease, Sjogren's syndrome and autoimmune hepatitis,¹⁴ although such an association has not always been found.^{15, 16}

A genome wide association study identified non-HLA genes associated with CD. After testing 310 605 single-nucleotide polymorphisms for association in 778 CD cases and 1422 controls, a linkage disequilibrium block encompassing the IL2 and IL21 genes was identified as a novel susceptibility factor.¹⁷ Additional genotyping recently identified seven previously unknown risk regions. Type-1 diabetes mellitus and CD shared several of these regions including HLA-DQ, IL2-IL21, CCR₃ and SH₂B₃.¹⁸ While these genetic associations will help understand pathogenic mechanism, they are not yet useful as predictors of the risk of developing severe complications of CD.

Of more potential use would be associations of particular HLA molecules with outcomes. HLA-DQ2 homozygosity has been associated with severe complications of CD; 26% of patients with refractory CD without aberrant T cells (RCD 1), 44% with refractory CD with aberrant T cells (RCD 2) and 54% with enteropathy-associated T-cell lymphoma (EATL) were homozygous for DQ2, compared with 21% with uncomplicated CD.¹⁹ The MYO9B gene on chromosome 19 also increases the risk for RCD 2 and EATL.²⁰ Homozygosity for the HLA-DQB1*0201 allele has been associated with a more severe clinical phenotype at presentation. In one report, it was associated with more marked villous atrophy, younger age, severe diarrhoea, a lower level of blood haemoglobin at diagnosis and a slower recovery of villous atrophy after a GFD.²¹ In another, HLA-DQB1*0201 homozygosity

Table 1. Prevalence of complications of and diseases associated with coeliac disease

Complication/ consequence	Details	Prevalence		Evidence for association (reference)			
		Disease in general population	Disease in CD	CD in disease	Probable	Possible	No
Autoimmune disease	Thyroid disease	10–20% women, 1–2% men ¹⁵⁶	0.3% (subclinical CD) ¹⁵⁷	1.86% ¹⁶³ 4.3% ¹⁶⁴ 5.4% ¹⁶⁵ 5.9% ¹⁶⁶	158–162, 164–166	167, 168	
	Type-1 diabetes mellitus	0.17% ¹⁶⁹ 0.5–1% ¹⁷⁰	4% ³⁴ 13% ¹⁵⁸ 14% ¹⁵⁹ 14% ¹⁶⁰ 26% ¹⁶¹ 41% (anti-thyroid antibodies) ¹⁶² 4% ³⁴ 6% ¹⁷¹ 7% (subclinical disease) ¹⁵⁷	4.4% ¹⁷² 4.6% ¹⁷³ 4.9% ¹⁷⁴ 5.7% ¹⁷⁵	34, 157, 170–172, 174–176		178 178
	Type-2 diabetes mellitus Latent autoimmune diabetes in adults (LADA)	7.2%–9% ¹⁷⁷ 0.3% ¹⁷⁹ 7–10% of adults with type-2 diabetes mellitus ^{179–181}	Unknown Unknown	Unknown Unknown			
	Autoimmune hepatitis	1.56% women ¹⁸⁶ 3–4% ¹⁸⁷ 13.3% ¹⁸⁸	1.1% ³⁴ 4% ¹⁸²	6.4% ¹⁸³ 16% ¹⁸⁴ 4.5% ¹⁸⁹	8, 34, 182, 183, 185 189	168	
	Sjogren's syndrome	0.09% ¹⁹⁴ 0.033% ¹⁹⁹ 0.071% ²⁰⁰ 0.159% ²⁰¹ 0.1% ²⁰³	0.1% ³⁴ 1.3% ³⁴	15% ¹⁹¹ 5.9% ¹⁹⁵	191 195, 196	190 192, 193 34, 197, 198 34, 202	
	Glomerulonephritis Microscopic colitis Addison's disease Systemic lupus erythematosus	0.5% ²⁰⁴ 7.3% ²⁰⁵	1.3% ³⁴	9% ²¹¹ 9.15% ⁸⁰ 9.3% ²¹²	80, 207–210	10, 34	
Liver abnormalities	Alopecia areata Hypertansamin asaemia		25% ²⁰⁶ 32% ²⁰⁷ 39% ²⁰⁸ 42% ²⁰⁹ 44% ²¹⁰				

Table 1. (Continued)

Complication/ consequence	Details	Prevalence		Evidence for association (reference)			
		Disease in general population	Disease in CD	CD in disease	Probable	Possible	No
Pancreatic disease	Primary biliary cirrhosis	0.012% ²¹³ 0.34% ²¹⁴	2.6% ²¹⁰ 3.7% ²¹⁵	0% ^{210, 216} 2% ²¹⁷ 2.7% ²¹⁸ 3.4% ²¹⁵ 3.5% ²¹⁹ 1.5% ²²²	215, 217, 219, 220	168	210, 216, 221
	Cryptogenic liver disease						
	Primary sclerosing cholangitis	0.014% ²²⁴		1.6% ²¹⁹		219, 220	
	Non-alcoholic fatty liver disease	10–24% ²²⁵		3.3% ²²⁶ 3.4% ²²⁷		220, 227	
Pancreatic disease	Elevated pancreatic enzymes		25% ²⁰⁶			206	
	Pancreatitis	0.004–0.2% ²²⁸ 11.5% ²²⁹		7% ²³⁰		230, 231	
Neuropsychiatric disease	Neurological abnormalities		6–10% ¹³⁴ 10% ²³² 12% ⁶⁵ 0.5% ²⁴⁶		65	65, 134, 232–241	Ataxia ²⁴² Epilepsy ²⁴³
	Psychosis	0.006% (adolescents) ²⁴⁴ 1% ⁶⁶				246, 247	248, 249
	Depression	0.2–2.2% ²⁴⁵ 7.06% ²⁵⁰ 9.5% ²⁵¹ 15.2% ²⁵² 2.2% ²⁵³	57% ²⁵⁴		254–257		
Reproductive function: male	Anxiety	4.8–6% ²⁵⁸ 10.6% ²⁵⁹ 18.1% ²⁵¹ 19.5% ²⁶⁰	71% ²⁵⁴		254		
	Sexual dysfunction	31% ²⁶¹ 52% ²⁶²			263, 264		
	Hypogonadism		7% ²⁶³				25, 263

Table 1. (Continued)

Complication/ consequence	Details	Prevalence		Evidence for association (reference)			
		Disease in general population	Disease in CD	CD in disease	Probable	Possible	No
Reproductive function: female	Sexual dysfunction	4.3% ²⁶¹			264		
	Menstrual abnormalities	5% (early menopause) ²⁶⁵ 3–7.6% (secondary amenorrhoea) ²⁶⁶	39% (amenorrhoea) ²⁶⁷		267, 268	269	
Bone disease	Infertility and obstetric problems	7.4% (infertility) ²⁷⁰ 10% (small for gestational age) ²⁷¹	9% ²⁶⁷	4.1% ²⁷² 8% ²⁷³	267, 268, 273–280		281–284
	Reduced bone mass	27.3% ²⁸⁵ 28–47% (men), 15–33% (women) ²⁸⁶	31–35% ⁸⁶ 75% ⁵⁰	1.9% ²⁸⁷	50, 86, 287–290		
Hyposplenism	Osteoporosis	3–6% (men), 1–4% (women) ²⁸⁶ 4.1% ²⁸⁵	7–14% ¹⁴⁰ 26% ⁸⁶ 47% (women), 50% (men) ⁸⁷	9.4% ²⁹²	86, 87, 140, 292		
	Fractures	0.9–8.7% ²⁹¹ 8% ²⁹³ 3.3% ²⁹⁴ 3.4% ²⁹⁵	9% ²⁹⁶ 25% ²⁹³ 35% ²⁹⁴ 44% ²⁹⁵ 20–47% ²⁹⁷ 16% ³⁰¹ 32.8% ³⁰² 77% ³⁰³	295–299	301, 303, 304	305	294, 300
Infection	Functional hyposplenism and splenic atrophy						
	Urinary tract infection Bacterial sepsis Tuberculosis	5% ³⁰⁹	17% ³⁰⁹		310	306, 307 308 311	
Immune deficiency	IgA deficiency	0.11% ³¹²	2.3% ³¹³ 2.6% ³¹⁴		313, 314		
	Pericarditis, myocarditis, cardiomyopathy			2.1%–5.8% ^{315–317} 4.4% ³¹⁸		315–318	319
Venous thromboembolism	Venous	1.4% ³²⁰	2.6% ³²⁰			320	
	Iron deficiency Low folate Low vitamin B6 Hyper-homocysteinaemia	1–1.1% ³²¹	46% ¹⁵⁷ 37% 20% ¹²⁷ 20% ⁸⁵	2.3–20% ^{322–326}	157, 322, 324–327		

Table 1. (Continued)

Complication/ consequence	Details	Prevalence		Evidence for association (reference)			
		Disease in general population	Disease in CD	CD in disease	Probable	Possible	No
Malignancy	Lymphoma		SIR 2.2–5.1 ^{8, 31, 327–337} RR 42.7 ^{29, 30} RR 9.7–12.3 ^{29, 30}	0.92% ³³⁵	8, 327–337		
Impaired QOL	Other				29–31, 328, 338, 339 340–343		

The strength of association was judged as 'probable' for reasonable case series where a clear association was found and judged as 'possible' for case reports of for case series in which only a weak association was found.

CD, coeliac disease; SIR, standardized incidence ratio; RR, relative risk.

was associated with severe intestinal damage (Marsh 3c) in a dose-dependent manner, but not with the clinical presentation of the disease.²² In contrast, the zygosity of the DQA1*0501 allele does not have a significant impact on the severity of CD.²¹ Thus, identification of DQB1*02 homozygous patients may help identify CD patients at risk for developing these severe complications.

Environmental

The major environmental factor associated with the risk of developing coeliac-related complications is persistent exposure to dietary gluten and this is discussed in detail below. It is reasonable to assume that known risk factors for those complications in the general population would also apply to patients with CD, but the validity of that extrapolation has not been formally tested. Examples include smoking as a risk factor for Graves' ophthalmology²³ and osteoporosis;²⁴ alcohol excess for early menopause in women and loss of gonadal function in men²⁵ resulting in osteoporosis²⁶ and transient postpartum thyroiditis for chronic thyroiditis.^{27, 28}

Gluten

Gluten provides the antigenic drive to maintain intestinal disease and its removal from the diet not only results in symptomatic improvement but also allows a reduction in small intestinal inflammation and permits mucosal healing with improved absorptive function. It is anticipated, therefore, that gluten exposure would be associated with increased risk and/or a worse course of complications. Two key questions arise.

The first is whether GFD alters the course of associated illnesses. When complications or associated conditions are already present at the initiation of a GFD, there is abundant evidence of improved course of many, but not all conditions. The published data are outlined in Table 3.

The second key question is whether compliance to a GFD reduces the risk of developing complications or associated illnesses. Acquiring such data is extraordinarily difficult. The strongest evidence for a preventive effect of GFD is on malignancy, particularly lymphoma. Patients with untreated CD have higher rates of malignancy than the general population [oropharyngeal and oesophageal cancer relative risk (RR) 22.7, *P* < 0.001; lymphoma RR 77.8,

Complication	Risk factor		
	Gluten exposure	Intestinal inflammation – mucosal injury	Nutritional deficiency
Autoimmune thyroid disease	Yes ^{34, 158} No ^{35, 36, 161}	Yes ³⁹ Possible ^{77, 78}	Possible ^{88, 89}
Type-1 diabetes mellitus	Yes ³⁴ No ^{35, 36, 344–347}	Possible ^{77, 78}	
Liver disease	Yes ^{34, 207–209, 218, 223, 227, 348} No ³⁵	Possible ^{77, 78, 82}	
Neurological disease	Yes ²⁴¹ No ³⁴⁹	Possible ^{65, 350}	Possible ^{83, 90}
Psychiatric disease	Yes ^{68, 254, 256} No (depression) ²⁵⁴	Possible ⁶⁷	
Impaired reproduction and fertility	Yes ^{263, 264, 268, 269, 274–279, 283, 351}		Possible ^{85, 274}
Bone disease	Yes ^{47, 53, 56–60, 86, 87, 293, 352} No ^{140, 289}	Yes ^{50–52, 86}	Yes ^{47, 85–87} No ⁵⁷
Hyposplenism	Yes – functional ³⁵³ No – atrophy ³⁵⁴	Yes ⁷³	
Infection		Yes ⁶³	
Cardiac disease	Yes ^{318, 355, 356}	Yes ³²⁰	Yes ⁸⁵
Thromboses			
Nutritional deficiencies	Yes ^{7, 51} No ¹²⁷		
Refractory coeliac disease	Possible ³⁹	Yes ^{72, 73}	
Malignancy	Yes ^{29–31, 337}	Yes ⁷³	

Table 2. The reported contribution of ongoing gluten exposure, intestinal inflammation and injury and nutritional deficiency to the risk of complications in patients with coeliac disease

$P < 0.001$;^{29, 30} and small bowel carcinoma standardized incidence ratio 25].³¹ In comparison, CD patients on a GFD for >5 years have the same overall risk of malignancy as the general population.^{29, 30} The mean age at diagnosis of CD for patients diagnosed with cancer was significantly higher than the age at diagnosis of CD in patients who did not develop a malignancy.³¹ Assuming that the patients diagnosed with CD at an older age had a longer duration of untreated CD, the results further support that the GFD is protective against the development of malignancy.

In a cohort of 4633 individuals, an increase in mortality, particularly because of cancer, was associated with the presence of elevated IgA tissue transglutaminase (tTG) antibodies.³² It was unclear whether this was because of undiagnosed and, therefore, untreated CD, or whether the tTG antibodies were a marker of serious diseases such as cancer or heart failure. Analysis of blood samples obtained from US military recruits between 1948 and 1954 reveal that 0.2% were seropositive for tTG and endomysial (EM) antibodies.³³

These cases with unrecognized serologically detected CD had a nearly fourfold increased risk of death compared with seronegative subjects.

There is currently conflicting evidence regarding GFD altering the risk of developing autoimmune disease. One study showed that the duration of exposure to gluten may predispose to autoimmune disorders,³⁴ whilst others have been unable to demonstrate this association.^{35, 36} One of the proposed mechanisms by which autoimmunity might develop, molecular mimicry, does implicate continuing exposure to gluten. Gliadin may activate T cells and allow for expansion of the B cells specific for autoantigens complexed with gliadin peptides and for subsequent autoantibody production. Such inflammatory responses may have the capacity to exist in genetically susceptible hosts and lead to chronic organ-specific autoimmune disease via epitope spreading.³⁷ More recently, however, it has been demonstrated that tTG knockout mice immunized with transglutaminase do not develop any pathology, whereas immunized wild type did develop periductal lymphocytic infiltrates in

Table 3. Effect of gluten-free diet on the course of associated illnesses

System	Evidence
Endocrine disorders	Normalization of subclinical hypothyroidism ¹⁵⁸ Improved glycaemic control and increased body mass index in children with type-1 diabetes mellitus ³⁵⁷ No differences in height, BMI, HbA1C, insulin dose and haemoglobin between type-1 diabetes mellitus with or without CD at diagnosis of CD ³⁴⁷
Liver disease	Normalization of isolated mild elevations in transaminase activity and nonspecific hepatic histological changes ^{34, 207, 209} Prevention of progression to hepatic failure in severe liver disease ($n = 5$) ^{223, 348}
Neurological disease	No improvement in peripheral neuropathy and/or autonomic dysfunction ($n = 32$) ³⁴⁹ Significant improvements in muscle strength and neurophysiological abnormalities in motor neuropathy ($n = 2$) ²⁴¹
Psychiatric disorders	Reduced proportion with anxiety from 74% vs. 26% ($P < 0.001$) ²⁵⁴ No change in the proportion of patients with depression (57% vs. 48%) – higher than in controls ²⁵⁴ Improvement in depressive symptoms 2–5 months after the commencement of a GFD – case reports ²⁵⁶ Increase in major serotonin and dopamine metabolite concentrations in brain – in parallel with improvements in symptoms ³⁵⁸
Fertility and reproduction	Increased sexual activity and satisfaction ^{263, 264, 268, 274, 275, 277–279, 283, 351} Reduced rates of infertility, miscarriage, intrauterine growth restriction and low birth weight ^{263, 264, 268, 274, 275, 277–279, 283, 351} Undiagnosed CD may be a risk factor for unfavourable foetal outcome ²⁷⁶
Cardiac disease	Improvement in myocarditis and dilated cardiomyopathy ^{318, 355, 356}
Hyposplenism	Functional hyposplenism may be reversible ³⁵³ Splenic atrophy usually irreversible ³⁵⁴
Malignancy	Remission in some patients with type-1 RCD ³⁹ Increased overall mortality and rate of malignancy with ongoing gluten ingestion ³³⁸ .
Bone disease	Improvement in bone mineral density ^{53–60} , including postmenopausal women and those without complete duodenal mucosal recovery ⁵³ Normal bone mineralization long term in children and adults ^{59, 352}
Nutritional status	Increase in body weight, body mass index, fat mass, bone mass and triceps skin fold thickness ³⁵⁹ Improved in fat mass but not indices of lean body mass ⁸⁴ Improved nutrition and height velocity in Asian children ³⁶⁰ Prevention of nutrient deficiencies, in particular, iron deficiency anaemia ^{7, 51} Improved calcium absorption from the gut ³⁶¹ though not to normal values ⁴⁷ Parathyroid hormone levels return to normal ^{47, 59} .
Quality of life	Improvement (limited data) ^{340–343} Benefit in 'silent' CD, with improvements in fatigue and general well-being ³⁴¹

GFD, gluten-free diet; RCD, refractory coeliac disease; CD, coeliac disease.

lacrimal glands.³⁸ This observation suggests that the development of autoimmunity against tTG can be a pathological event which may lead to organ damage.

Prevention of many of the other complications that are a direct result of intestinal and systemic inflammation and/or nutritional deficiencies, such as bone disease, is assumed as GFD is able to attenuate

and, in many cases, correct such pathogenic mechanisms. However, it is unknown whether adherence to a GFD reduces the chance of subsequently developing conditions in which the pathogenic mechanisms is uncertain (such as autoimmune disease, infertility, neuropsychiatric disorders and liver disease). Patients with CD must be considered as having an ongoing risk.

Ongoing mucosal inflammation and injury

If ongoing inflammation and injury in the intestinal mucosa are pathogenically involved in complications and associated conditions, then ensuring healing of the intestinal lesion becomes a key target in their prevention. The clearest evidence of the pathogenic importance of ongoing inflammation in the development of complications of CD derives from observations in patients with type-1 RCD, where intestinal inflammation persists independently of gluten ingestion. These patients are at increased risk of concomitant autoimmune diseases and infectious and thromboembolic complications.^{39, 40} In fact, despite clinical improvement on the GFD, many histologically nonresponsive CD patients develop severe complications, such as gastrointestinal cancer and lymphoma, within a relatively short follow-up.⁴⁰

Several mechanisms by which intestinal immune and inflammatory events may contribute to systemic symptoms and development of other conditions have been implicated. First, the production of cytokines and other inflammatory mediators as an integral part of immune activation at the mucosal level is associated with systemic exposure to such factors. Circulating levels of proinflammatory cytokines are increased in patients with untreated CD including IL-1 β , IL-2, IL-2 receptor, IL-6, IL-10, IL-18, IFN- γ , TNF- β/α indicative of T-cell activation,⁴¹⁻⁴⁵ and reduction in levels of anti-inflammatory cytokines, such as IL-1 receptor antagonist, has also been observed.⁴¹ Of importance, circulating levels of proinflammatory and anti-inflammatory cytokines significantly decrease and increase, respectively, on effective treatment of CD with a GFD.^{41, 42} These manifestations of 'systemic inflammation' are likely to contribute to major symptoms and signs of untreated CD, such as fatigue and weight loss. Cytokines increase bone turnover and induce cortical bone loss via direct effects on osteoclastogenesis and osteoblast activity and these have been demonstrated in patients with CD.⁴⁶⁻⁵⁰ Such a mechanism probably underlies the greater reduction in BMD in patients with villous atrophy compared with those with less histologic severity.^{51, 52} Treatment with GFD and the subsequent reduction in systemic inflammation are associated with improved BMD⁵³⁻⁶⁰ as outlined above.

Secondly, systemic cellular changes occur and these may be responsible for the increased risk of infection. Untreated patients have significantly reduced levels of leucocytes, total lymphocytes, and CD3+, CD4+ and

CD8+ lymphocytes compared to those with treated CD and to the general population.⁶¹⁻⁶³ Hyposplenism/splenic sequestration (see Table 1) is one mechanism. The proportion of natural killer and cytotoxic T cells is lower in untreated and treated coeliac patients than in healthy populations, postulated to be because of sequestration of lymphocytes within the intestinal mucosa and/or because of their loss into the gut lumen.⁶³ Furthermore, excessive apoptosis of T cells has been reported in CD and this may contribute to the observed lymphopenia.⁶⁴

Thirdly, shared immunological abnormalities suggest a common underlying vulnerability between active CD and neuropsychiatric disorders.^{65, 66} It has been postulated that gluten may exert its pathogenic effect on processes in the brain through cell-mediated immune mechanisms and lymphocyte stimulation.⁶⁷ Impaired availability of tryptophan and disturbances in central serotonergic function may play a role in the development of anxiety, headaches/migraines, behavioural symptoms and depression in CD.^{68, 69} These disturbances may be caused by interferon- γ , the predominant cytokine produced by gluten-specific T cells in active CD.⁷⁰ Interferon- γ can suppress serotonin function both directly and indirectly by enhancing serotonin turnover.⁷¹

Fourthly, immunological events can lead to local intestinal complications. In patients with type-2 RCD, clonal expansion of intraepithelial lymphocytes appears to be driven by IL-15 secreted by epithelial cells. This causes proliferation of intraepithelial lymphocytes independent of gluten stimulation. This results in the secretion of interferon- γ , which is cytotoxic to intestinal epithelial cells.⁷² Patients with this type of RCD are at high risk of developing ulcerative jejunitis, mesenteric lymph node cavitation, hyposplenism and small intestinal lymphoma⁷³ and have a poor prognosis.⁷⁴

Finally, mucosal inflammation is associated with increased intestinal permeability, because of epithelial injury⁷⁵ and possibly altered zonulin signalling.⁷⁶ Such increased leakiness of the epithelium has been implicated in several aspects of the symptomatology and complications of CD. It may predispose to other autoimmune disorders by facilitating further external antigens such as food proteins, bacterial products and endotoxins to enter the intestinal lamina propria, thus leading to the activation of autoimmune phenomena.⁷⁷⁻⁷⁹ Entry of these molecules into the portal circulation has been postulated to cause

hypertransaminasaemia^{80, 81}, which is significantly associated with increased intestinal permeability in patients with CD.⁸² Likewise, systemic exposure to xenobiotics may contribute to tiredness and other nonspecific symptoms. It has also been implicated in the pathogenesis of acquired neurological disorders,⁸³ but the nature of the putative toxins remains unknown.

Nutritional deficiencies

It is important to recognize nutritional deficiencies that impact upon the likelihood of complications of CD occurring as these can usually be readily corrected. There is relatively little information of the influence of macronutrient malnutrition, except in children where this is likely to be associated with developmental delay and/or growth failure. In adults, it is unusual for indices of lean body mass to be deficient amongst patients diagnosed with CD today.⁸⁴ In contrast, micronutrient deficiencies are well recognized and relatively common, as follows:

Iron deficiency may lead to anaemia, tiredness and cognitive impairment.

Folate deficiency may also contribute to anaemia, but is also with associated elevated homocysteine levels. Hyper-homocysteinaemia is a risk factor for thrombosis, recurrent miscarriages and osteoporosis.⁸⁵

Intestinal malabsorption of *calcium* and *vitamin D* may also contribute to osteoporosis and osteomalacia.^{47, 86, 87}

The body content of *selenium* may be low in untreated CD⁸⁸ and this may impair the action of thyroid hormones,⁸⁹ although there are currently no data on the association between thyroid function and selenium deficiency in patients with untreated CD.

Vitamin E deficiency has been postulated to be one of the causes of neurological abnormalities, in particular, cerebellar disease.^{83, 90} Other nutrient deficiencies have been postulated without evidence.⁸³

METHODS OF EVALUATING RISK FACTORS

Assessing genetic risk factors

The utility of genetic testing to estimate risk in CD is still in its infancy and should not be applied to routine clinical practice at present. However, there are suggestions that gene dosage, subtypes and zygosity of HLA-DQ2 and DQ8 may be useful predictors of which of the patients should be closely followed in the

future.^{21, 22, 91} Other genes associated with CD will need to be assessed in this context.

Assessing environmental risk factors

At each review, environmental risk factors independent of gluten and inflammation need addressing. Important factors include smoking, alcohol excess and obesity.

Monitoring adherence to the GFD

Assessment of compliance to the GFD is challenging. There are no clear guidelines as to the optimal method for monitoring adherence to this diet. However, a multidisciplinary assessment approach is needed:

Dietary history. Dietary compliance as assessed by a skilled dietitian trained in GFD at interview is thought to be one of the best markers of dietary adherence because of low cost, non-invasiveness and correlation with intestinal damage.⁹² Adherence to the diet requires a focused dietary history and it is essential that dietitians are up-to-date for the current teaching of a GFD.³

Clinical assessment. Lack of strict adherence to the GFD is the most common cause of persistent symptoms in CD.^{92, 93} However, resolution of symptoms may not be accurate in assessing adherence to the GFD as judged by interview and biopsy.⁹⁴ Symptoms may also indicate issues other than gluten ingestion; for example, malabsorption of fructose and lactose is common in untreated CD, irritable bowel symptoms may worsen with the GFD because more fruits or 'naturally' gluten-free foods rich in fructose are consumed⁹⁵ and occasionally symptoms represent the development of complications such as RCD or lymphoma.

Coeliac serology. Levels/titres of coeliac-associated antibodies fall with reduction in gluten intake. EM antibody and tTG IgA levels have been shown to normalize in seven out of eight patients within a year on a strict GFD.⁹⁶ However, the rate of fall, speed of its normalisation or whether serology normalizes or not are not reliable markers of strict adherence to the GFD⁹⁷⁻⁹⁹ and cannot replace expert nutritionist assessment.⁹⁹

Likewise, normalization of the levels of, for example, that of EM antibody levels, occurs in 87% of patients after 12 months on a GFD, but this is an unreliable marker of healing of the intestinal lesion^{96, 97, 100–102} arguably a good index of strict adherence (see below). After normalization, serological markers can then be used as indicators of ongoing gluten-free dietary compliance.^{96, 103, 104} Thus, serial serology has greater utility in predicting non-adherence than strict adherence.

Duodenal biopsy. Small bowel histological assessment is the gold standard for assessing strict compliance, as indicated by healing of the mucosa. To complicate the issue, some patients who have healed their intestinal mucosa can ingest gluten without inducing further histological abnormality.^{105, 106} Interpretation of the histopathology with respect to gluten intake must also be made in light of previous biopsy appearances when taking a GFD. This is a strong argument for performing a follow-up biopsy after 12 months or so of apparent strict GFD.

Assessing intestinal inflammation and mucosal injury

The definitive way of assessing the degree of healing and mucosal inflammation is via the assessment of small bowel histology. A repeat biopsy on the GFD will help define mucosal healing and identify persistent mucosal injury, even in the absence of symptoms or positive serology. Despite apparent adherence to the GFD, histological recovery has been demonstrated to take time; for example, 62% of 57 patients had persisting villous atrophy over a median of 13 months⁸⁴ and in other cohorts, 40% had incomplete healing after 24 months¹⁰⁷ and 10% after 5 years.¹⁰⁸ The criteria for diagnosis of CD from the European Society of Paediatric Gastroenterology and Nutrition do not demand follow-up biopsy,¹⁰⁹ but, given the high rate of persistent villous atrophy despite the GFD, a repeat biopsy should be considered 1–2 years after its commencement.¹⁰⁸

A minimum of four endoscopic forceps biopsies of the descending duodenum are known to enhance the value of small bowel histology.^{110, 111} Thus, obtaining only two biopsy specimens will lead to a confirmed diagnosis in only 90%, whereas four specimens will lead to 100% confidence in diagnosis.¹¹¹ However,

because of the patchy nature of villous atrophy, a five-biopsy regimen including one from the duodenal bulb, two from the first part and two from the second part of the duodenum, is required for detection of the most severe lesion.¹¹² The issue of patchiness of the lesions is also being addressed by techniques such as magnification endoscopy,^{113, 114} confocal laser microscopy^{115, 116} or capsule endoscopy,¹¹⁷ although optimal roles for these technologies in clinical practice are yet to be clearly established.

Non-invasive ways of assessing healing and inflammatory activity are desirable so that they can be repeated at more frequent intervals. Potential techniques include the following:

Clinical assessment. Despite clinical improvement on the GFD, villous atrophy may persist.^{40, 97, 118, 119} Symptom assessment cannot be advocated as a reliable method for determining absence of persistent small bowel villous atrophy.

Coeliac serology. Normalization of coeliac antibodies does not in itself guarantee that the small intestinal villous structure has improved.^{40, 97, 100} For example, in a series of 57 consecutive patients, 53% had normalized tTG IgA, but three out of five of these had ongoing villous atrophy. Conversely, one-third of the patients who still had raised tTG antibody levels at 12 months had normal histology.⁸⁴ In a series of 13 patients with persistent villous atrophy and crypt hyperplasia, tTG IgA was normal in 59% and EM antibody normal in 74%.⁴⁰ However, persistently positive serology or rising titres may indicate mucosal injury or the development of RCD.¹²⁰

Markers of systemic inflammation. The use of circulating cytokine levels has not progressed from a research tool to clinical practice. The utility of commonly used markers such as the erythrocyte sedimentation rate or levels of C-reactive protein has not been reported.

Markers of absorptive surface area and function. No marker of absorptive function has proven utility in defining small bowel surface area and therefore the degree of villous atrophy. Candidates include the ¹³C-sucrose breath test, which has been a useful and

sensitive biomarker for defining changes in the severity of mucositis in patients receiving cancer chemotherapy,^{121, 122} or plasma citrulline levels, which correlate with small bowel length and xylose absorption.¹²³

Intestinal paracellular permeability. Dual sugar permeability studies are frequently abnormal in CD,^{124, 125} improve rapidly on the GFD¹²⁶ and appear to be sensitive in detecting even small amounts of gluten in the diet.¹²⁵ However, their use in longitudinally evaluating patients with CD has not been reported.

Testing for nutritional deficiencies

Fortunately, the key nutritional markers outlined above can be measured in routine laboratories and methods for their correction are readily available. Despite adherence to the GFD, nutritional deficiencies may still exist. A Swedish study showed that despite a GFD for several years, half of adult coeliac patients still showed signs of a poor vitamin status with elevated homocysteine levels, low folate and low vitamin B6 levels.¹²⁷ This may be attributed, in part, to reduced dietary intake of nutrients because of the strict GFD. The results suggest that when following up patients with CD, the micronutrient status should be reviewed even in patients who have a good response to the GFD.

RECOMMENDATIONS FOR FOLLOW-UP

There are four key issues that require addressing:

Defining whether follow-up is indicated

Given the high number of complications associated with CD, the discovery of several risk factors, which may cause these or worsen outcome and the development of ways to assess risk, it appears intuitive that follow-up is necessary. Gluten ingestion is one of the key risk factors for developing complications and regular follow-up and improved knowledge in both children and adults have been shown to correlate positively with dietary compliance.^{128–131} The concept that the intensity of follow-up can be stratified according to phenotypic characteristics of the individual patients has little supportive evidence. Symptoms and serology are poor guides to the severity of the intestinal lesion and to nutritional

and bony status. The prevalence of persistent histological changes on the GFD is high, being up to 60% at a median of 2 years.¹¹⁹ As this is a clear risk factor for complications,⁴⁰ follow-up is indicated for all patients.

Defining who should administer follow-up

There is a lack of evidence on who should follow-up patients with CD, whether a specialist gastroenterologist, general physician, general practitioner or dietitian. Instinctively, someone who has both knowledge and interest in CD would be optimal. When surveyed, patients with CD would prefer to see a dietitian for follow-up with a doctor being available.¹³² Provided that the dietitian was trained in assessing and instigating the GFD and that clear guidelines on when to call for medical help are in place, this approach would allow good assessment of dietary compliance³ and offer opportunities for reinforcement of dietary principles, with less medical input for a majority of coeliac patients who remain well. A doctor trained in the management of CD would be required at initial consultation for those patients who require more intensive management and to monitor laboratory investigations to detect nutritional deficiencies or other complications.

Defining what should be monitored and how often

Recommendations on the frequency of follow-up cannot be made on a solid evidence-base, but derive from what seems reasonable. Following histological diagnosis of CD, the minimum requirements for all patients are an initial consultation 1–2 weeks after endoscopy, review consultation 3–6 months later and subsequent reviews yearly,¹³³ although this would need to be individually assessed. A suggested plan of management is illustrated in Figure 1.

Follow-up should address several aspects:

Compliance to GFD. The central platform of gaining optimal outcomes in patients with CD is to ensure that they adhere to the GFD. At each review, dietary compliance should be assessed by history and serology. A dietitian is best equipped to evaluate by dietary history and to correct any misconceptions and educate regarding inadvertent gluten ingestion. As




Figure 1. A suggested plan of follow-up management of patients with coeliac disease. ^aCoeliac serology, FBC, electrolytes, LFTs, thyroid function, iron studies, calcium, phosphate, vitamin D, folate, B₁₂, fasting glucose, ± zinc, Mg. ^bAt 1–2 years; then as indicated on clinical grounds. ^c3–5 yearly in high-risk groups (noncompliance, refractory coeliac disease, female ≥50 years, fractures, men ≥55 years); yearly if osteoporosis on treatment.

outlined above, the results of serological tests, tTG IgA and EM antibodies, have to be interpreted carefully to avoid false security or inappropriate anxiety in the patient. For the medical practitioner, every

review should be an opportunity to promote compliance. Ensuring membership of a patient-led self-help organisation is essential as these organisations provide detailed dietary information, food lists and

educational material in addition to social support.¹²⁹ Neuropsychiatric disturbances, such as depression, may adversely affect compliance¹³⁴ and must be addressed and managed. Anger can worsen the patient-clinician relationship and has been inversely correlated with dietary compliance.¹³⁵

Symptoms. At each review, a thorough history of gastrointestinal symptoms and other symptoms associated with related conditions and complications such as neuropsychiatric symptoms, fatigue and sexual dysfunction may guide future management.

Duodenal biopsy. The need for repeated assessment of duodenal histopathology has been controversial and practice in this way varies across practitioners. As outlined above, symptom response and serology are poor predictors of healing in the individual patient and accurate non-invasive methods of assessing healing are yet to be validated. Demonstrating that the intestinal lesion has healed provides reinforcement and reward for the patient's efforts, provides important information for the appropriate interpretation of biopsies that may be subsequently performed because of the appearance of gut symptoms or other issues and permits recognition of patients with RCD before serious complications may develop. The main disadvantage of performing follow-up biopsies is the risk of inducing anxiety in patients because of undue expectation of having achieved healing. Thus, appropriate counselling needs to be linked to the revelation of the results.

The appropriate timing of the follow-up biopsy after GFD is initiated has not been established, but the earlier it is, the higher the chance that healing will not have occurred. Re-biopsy no earlier than 1–2 years after commencement of the GFD is recommended.^{107, 108} If the mucosa has healed, further biopsies cannot be justified unless indicated on changes in clinical status. If histological improvement is incomplete, biopsies probably should be performed again in another 1–2 years. No improvement requires a different strategy, comprising at least dietary re-evaluation and repeat biopsies in a further 6–12 months. In some clinical scenarios, imaging of the small bowel maybe indicated (see discussion below).

Nutritional assessment. Micronutrient deficiencies are common even in patients who are adherent to the

GFD.¹²⁷ For this reason, the key markers of nutritional state should be evaluated at each review. Thus, full blood evaluation, iron studies, and the levels of vitamin D, calcium, phosphate, folate and B₁₂ should be measured. Other markers that could be considered in selected patients and/or on some occasions include: homocysteine levels, particularly in those patients with a history of venous thromboembolism; parathyroid hormone concentrations as an indicator of secondary hyperparathyroidism²⁴ as serum calcium, phosphate and alkaline phosphatase measurements may be normal despite the presence of osteomalacia; serum magnesium as this was low in 13% one population reported;¹³⁶ and serum zinc as this is reduced in up to one in three patients with CD.¹³⁶ Assessing the levels of vitamin E and selenium (as discussed above) must be considered optional. Deficiencies should be corrected by specific supplementation and subsequent monitoring schedules adjusted accordingly.

Screening for complications and associated diseases. As the new development of associated diseases/complications is relatively uncommon in patients with CD, any monitoring should be simple and inexpensive. Thyroid function tests, a random or fasting blood glucose, electrolytes, liver function tests and platelet count should be routinely performed at every review. Homocysteine levels should be monitored in the appropriate clinical situation (as discussed above).

Bone densitometry. The timing of bone densitometry is controversial. The British Society of Gastroenterology²⁴ recommends that all patients have bone densitometry performed at diagnosis and regularly thereafter on the basis of increased fracture risk.^{24, 137} This seems reasonable given the fact that reduced bone density cannot be predicted on the basis of symptoms, blood results or duodenal histology.⁸⁴ However, the actual risk of fracture in patients with CD is still unclear and the predictive value of bone densitometry is not sufficient to identify accurately individuals who will sustain fractures.^{137, 138} The recommended approach to perform densitometry regularly after the initial study is not widely supported and requires further studies on safety, efficacy and cost-effectiveness.^{139–141} Follow-up bone density assessment should probably be restricted to high-risk situations that include patients with osteopenia/porosis at diagnosis,

those with RCD, post-menopausal women, patients who have sustained minimal trauma fractures and perhaps men >55 years. The frequency of bone densitometry should be dictated by the scenario; for example, post-menopausal women should be screened every 3–5 years, whereas patients undergoing treatment for osteoporosis may require yearly assessments.²⁴

Defining a strategy for 'red flags' for intestinal complications

A schedule of routine monitoring requires strategies for dealing with abnormalities detected. Identification of specific micronutrient deficiencies or associated conditions such as thyroid disease or hyposplenism leads to clear plans of action that are beyond the scope of this review. However, 'red flags' for more serious intestinal complications of CD, namely RCD, ulcerative jejunoileitis and small bowel lymphoma, should be recognized and subsequently addressed.

Such red flags include the following: ongoing, increasing or newly developed gastrointestinal symptoms; the development of associated symptoms such as unexplained fever, weight loss and night sweats; ongoing or new nutritional deficiencies and increasing levels of coeliac antibodies. The strategy should include:

Dietary exposure to gluten should be re-evaluated by a specialist dietitian and more extensive counselling regarding the GFD should be given. There are two additional key issues to consider. First, the style of GFD that was taught should be defined. Guidelines on how gluten-free the diet should be vary throughout the world, with recommendations on permissible gluten ranging from 20 to 200 mg/day. For example, in the UK, the 'limited detectable gluten diet', which permits ingestion of minute amounts of gluten up to the limit recommended by the Codex Alimentarius, is recommended, whereas the 'no-detectable gluten' diet is applied in Australia and the 'zero-tolerance' GFD in Canada. One of the larger studies addressing the safety threshold of prolonged exposure to trace amounts of gluten recommended that ingestion of contaminating gluten should be kept lower than 50 mg/d in the treatment of CD.¹⁴² A recent systematic review of thirteen studies assessing permissible gluten levels found that while many patients tolerated an average of 34–36 mg of gluten per day, others who consumed about 10 mg of gluten per day developed mucosal abnormalities.¹⁴³ It is intuitive

that patients with incomplete histological recovery must be advised to ensure as close to complete exclusion of gluten from their diet.

Secondly, the consumption of oats must be considered. Whether oats is allowed in the GFD is a contentious issue and opinions are divided on toxicity of oats in CD.^{144–147} While oats may be tolerated by most coeliac patients, some patients are sensitive to oats¹⁴⁴ and commercially available oats are often contaminated by wheat, barley and rye.¹⁴⁸ It seems only sensible that patients who do not have their intestinal lesion healed should avoid oats until clinical and histological remission is achieved.^{144, 146} Subsequently, up to 50–70 g/day of noncontaminated oats may be added.^{145, 146}

Alternative causes should be contemplated. For example, gut symptoms should prompt consideration of the presence of irritable bowel syndrome, itself often amenable to additional dietary changes¹⁴⁹ or other causes of duodenal villous atrophy and malabsorption such as autoimmune enteropathy, inflammatory bowel disease, common variable immune deficiency syndrome, tropical sprue and eosinophilic gastroenteritis should be reconsidered; recurrent iron deficiency should prompt consideration of colorectal neoplasia in the older patient or excessive menstrual blood loss in young women.

A repeat duodenal biopsy should be seriously considered after ensuring adequate dietary compliance. This provides not only useful information for both the patients and the attending doctor but also the key investigation in identifying patients at risk of proceeding to severe clinical manifestations such as RCD, ulcerative jejunoileitis and small bowel lymphoma.⁴⁰

Special investigations may then be necessary, if the duodenal mucosa is atrophic or lymphoma is suspected. Immunophenotyping or immunohistochemistry of the intestinal mucosa can provide important clues. They may demonstrate the characteristic expanded intraepithelial lymphocyte (IEL) population that lacks CD4, CD8 and T-cell receptor β -chain¹⁵⁰ and may be useful in confirming the diagnosis of RCD. Aberrant cells of more than 20% are considered abnormal. Clonality assessment for TCR- γ gene rearrangements is achieved using a TCR-PCR protocol. A clonal T-cell population can be found in RCD and is thought to be predictive of the development of EATL.^{39, 151} Increased lactate dehydrogenase or β 2-microglobulin is a strong indicator of lymphoma in addition to clonal TCR gene

rearrangements.¹⁵² In patients on a GFD, EATL may not be accompanied by duodenal villous atrophy. If an overt lymphoma is suspected, an extensive work-up is required including gastroscopy with multiple duodenal biopsies, CT chest, abdomen and pelvis.⁷⁴ Other potentially useful investigations include CT or MR enteroclysis,¹⁵³ PET scanning,¹⁵⁴ video capsule enteroscopy and double balloon enteroscopy.^{74, 155} In some cases, laparotomy, intraoperative enteroscopy and full thickness biopsies are necessary.

CONCLUSIONS

The bulk of evidence suggests that CD is not a trivial disease and requires not only early diagnosis but also planned, long-term follow-up. While a vast majority of patients will have an excellent long-term outlook, current understanding does not permit stratification of patients for various levels of intensive review. The presence of symptoms is no guide to the severity of intestinal, bony or nutritional abnormalities and should not be the principal guide to the initial or long-term management strategy. Improved methods to identify those patients at high risk of developing complications and/or associated diseases are needed. Duodenal biopsy remains the

best means of checking dietary compliance and detecting ongoing intestinal inflammation and villous atrophy, but it remains an invasive and expensive test that cannot be frequently repeated. Reliable non-invasive methods of assessing the status of the duodenal mucosa will greatly assist in more precise follow-up.

ACKNOWLEDGEMENTS

Declaration of personal interests: (i) Robert Anderson has served as a speaker, a consultant and an advisory board member for AstraZeneca Australia, Given Imaging, BTG International Plc and Alvine Pharmaceuticals. Peter Gibson has served as a consultant for Alvine Pharmaceuticals. (ii) Robert Anderson is a substantial shareholder, Director and Chief Scientist/Medical Officer of Nexpep Pty Ltd and Nexgrain Pty Ltd. (iii) Robert Anderson is named inventor of patents PCT/GB00/03760, PCT/GB03/002450 and 2004201774 pertaining to the diagnostic, therapeutic and functional food applications of T-cell epitopes in CD. *Declaration of funding interests:* Robert Anderson has received research funding from NHMRC, CoeliacUK, BTG International Plc, Ramaciotti Foundation, CSIRO/GRDC and Nexpep Pty Ltd.

REFERENCES

- Anderson RP. Coeliac disease: current approach and future prospects. *Int Med J* 2008; in press.
- Lohi S, Mustalahti K, Kaukinen K, *et al*. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther* 2007; **26**: 1217–25.
- Shepherd S, Gibson PR. Understanding the gluten-free diet for teaching in Australia. *Nutr Diet* 2006; **63**: 155–65.
- Polvi A, Eland C, Koskimies S, *et al*. HLA DQ and DP in Finnish families with celiac disease. *Eur J Immunogenet* 1996; **23**: 221–34.
- Michalski JP, McCombs CC, Arai T, *et al*. HLA-DR, DQ genotypes of celiac disease patients and healthy subjects from the West of Ireland. *Tissue Antigens* 1996; **47**: 127–33.
- Polvi A, Arranz E, Fernandez-Arquero M, *et al*. HLA-DQ2-negative celiac disease in Finland and Spain. *Hum Immunol* 1998; **59**: 169–75.
- Rostom A, Dube C, Cranney A, *et al*. Celiac disease. *Evid Rep Technol Assess (Summ)* 2004; **104**: 1–6.
- Rostom A, Murray JA, Kagnoff MF. American Gastroenterological Association (AGA) Institute technical review on the diagnosis and management of celiac disease. *Gastroenterology* 2006; **131**: 1981–2002.
- Atkinson MA, Eisenbarth GS. Type 1 diabetes: new perspectives on disease pathogenesis and treatment. *Lancet* 2001; **358**: 221–9.
- Kaukinen K, Collin P, Mykkanen AH, *et al*. Celiac disease and autoimmune endocrinologic disorders. *Dig Dis Sci* 1999; **44**: 1428–33.
- Wallaschofski H, Meyer A, Tuschy U, *et al*. HLA-DQA1*0301-associated susceptibility for autoimmune polyglandular syndrome type II and III. *Horm Metab Res* 2003; **35**: 120–4.
- Segni M, Pani MA, Pasquino AM, *et al*. Familial clustering of juvenile thyroid autoimmunity: higher risk is conferred by human leukocyte antigen DR3-DQ2 and thyroid peroxidase antibody status in fathers. *J Clin Endocrinol Metab* 2002; **87**: 3779–82.
- Morling N, Dalhoff K, Fugger L, *et al*. DNA polymorphism of HLA class II genes in primary biliary cirrhosis. *Immunogenetics* 1992; **35**: 112–6.
- Dalton TA, Bennett JC. Autoimmune disease and the major histocompatibility complex: therapeutic implications. *Am J Med* 1992; **92**: 183–8.
- Taylor JC, Gough SC, Hunt PJ, *et al*. A genome-wide screen in 1119 relative pairs with autoimmune thyroid disease. *J Clin Endocrinol Metab* 2006; **91**: 646–53.
- Gillett HR, Cauch-Dudek K, Jenny E, *et al*. Prevalence of IgA antibodies to endomysium and tissue transglutaminase

- in primary biliary cirrhosis. *Can J Gastroenterol* 2000; 14: 672–5.
- 17 van Heel DA, Franke L, Hunt KA, *et al.* A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. *Nat Genet* 2007; 39: 827–9.
- 18 Hunt KA, Zernakova A, Turner G, *et al.* Newly identified genetic risk variants for celiac disease related to the immune response. *Nat Genet* 2008; 40: 395–402.
- 19 Al-Toma A, Goerres MS, Meijer JW, *et al.* Human leukocyte antigen-DQ2 homozygosity and the development of refractory celiac disease and enteropathy-associated T-cell lymphoma. *Clin Gastroenterol Hepatol* 2006; 4: 315–9.
- 20 Wolters VM, Verbeek WH, Zernakova A, *et al.* The MYO9B gene is a strong risk factor for developing refractory celiac disease. *Clin Gastroenterol Hepatol* 2007; 5: 1399–405, 405 e1–2.
- 21 Karinen H, Karkkainen P, Pihlajamaki J, *et al.* Gene dose effect of the DQB1*0201 allele contributes to severity of coeliac disease. *Scand J Gastroenterol* 2006; 41: 191–9.
- 22 Jores RD, Frau F, Cucca F, *et al.* HLA-DQB1*0201 homozygosity predisposes to severe intestinal damage in celiac disease. *Scand J Gastroenterol* 2007; 42: 48–53.
- 23 Bartalena L, Martino E, Marcocci C, *et al.* More on smoking habits and Graves' ophthalmopathy. *J Endocrinol Invest* 1989; 12: 733–7.
- 24 Scott EM, Gaywood I, Scott BB. Guidelines for osteoporosis in coeliac disease and inflammatory bowel disease. British Society of Gastroenterology. *Gut* 2000; 46(Suppl. 1): i1–8.
- 25 Farthing MJ, Rees LH, Edwards CR, *et al.* Male gonadal function in coeliac disease: 2. Sex hormones. *Gut* 1983; 24: 127–35.
- 26 Finkelstein JS, Klibanski A, Neer RM, *et al.* Osteoporosis in men with idiopathic hypogonadotropic hypogonadism. *Ann Intern Med* 1987; 106: 354–61.
- 27 Othman S, Phillips DI, Parkes AB, *et al.* A long-term follow-up of postpartum thyroiditis. *Clin Endocrinol (Oxf)* 1990; 32: 559–64.
- 28 Jansson R. Postpartum thyroid disease. *Mol Biol Med* 1986; 3: 201–11.
- 29 Cooper BT, Holmes GK, Cooke WT. Lymphoma risk in coeliac disease of later life. *Digestion* 1982; 23: 89–92.
- 30 Holmes GK, Prior P, Lane MR, *et al.* Malignancy in coeliac disease – effect of a gluten free diet. *Gut* 1989; 30: 333–8.
- 31 Silano M, Volta U, Mecchia AM, *et al.* Delayed diagnosis of coeliac disease increases cancer risk. *BMC Gastroenterol* 2007; 7: 8.
- 32 Metzger MH, Heier M, Maki M, *et al.* Mortality excess in individuals with elevated IgA anti-transglutaminase antibodies: the KORA/MONICA Augsburg cohort study 1989–1998. *Eur J Epidemiol* 2006; 21: 359–65.
- 33 Rubio-Tapia A, Lahr B, Zinsmeister AR, *et al.* *The Silence of the Intestines: Increased Prevalence and Mortality of Undiagnosed Celiac Disease.* San Diego, CA: DDW, 2008.
- 34 Ventura A, Magazzu G, Greco L. Duration of exposure to gluten and risk for autoimmune disorders in patients with celiac disease. SIGEP Study Group for Autoimmune Disorders in Celiac Disease. *Gastroenterology* 1999; 117: 297–303.
- 35 Viljamaa M, Kaukinen K, Huhtala H, *et al.* Coeliac disease, autoimmune diseases and gluten exposure. *Scand J Gastroenterol* 2005; 40: 437–43.
- 36 Sategna Guidetti C, Solerio E, Scaglione N, *et al.* Duration of gluten exposure in adult coeliac disease does not correlate with the risk for autoimmune disorders. *Gut* 2001; 49: 502–5.
- 37 Schuppan D. Current concepts of celiac disease pathogenesis. *Gastroenterology* 2000; 119: 234–42.
- 38 Freitag T, Schulze-Koops H, Niedobitek G, *et al.* The role of the immune response against tissue transglutaminase in the pathogenesis of coeliac disease. *Autoimmun Rev* 2004; 3: 13–20.
- 39 Al-Toma A, Verbeek WH, Mulder CJ. Update on the management of refractory coeliac disease. *J Gastrointest Liver Dis* 2007; 16: 57–63.
- 40 Kaukinen K, Peraaho M, Lindfors K, *et al.* Persistent small bowel mucosal villous atrophy without symptoms in coeliac disease. *Aliment Pharmacol Ther* 2007; 25: 1237–45.
- 41 Fornari MC, Pedreira S, Niveloni S, *et al.* Pre- and post-treatment serum levels of cytokines IL-1beta, IL-6, and IL-1 receptor antagonist in celiac disease. Are they related to the associated osteopenia? *Am J Gastroenterol* 1998; 93: 413–8.
- 42 Romaldini CC, Barbieri D, Okay TS, *et al.* Serum soluble interleukin-2 receptor, interleukin-6, and tumor necrosis factor-alpha levels in children with celiac disease: response to treatment. *J Pediatr Gastroenterol Nutr* 2002; 35: 513–7.
- 43 Lettesjo H, Hansson T, Bergqvist A, *et al.* Enhanced interleukin-18 levels in the peripheral blood of children with coeliac disease. *Clin Exp Immunol* 2005; 139: 138–43.
- 44 Penedo-Pita M, Peteiro-Cardelle J. Increased serum levels of interleukin-2 and soluble interleukin-2 receptor in celiac disease. *J Pediatr Gastroenterol Nutr* 1991; 12: 56–60.
- 45 Lahat N, Shapiro S, Karban A, *et al.* Cytokine profile in coeliac disease. *Scand J Immunol* 1999; 49: 441–6.
- 46 Selby PL, Davies M, Adams JE, *et al.* Bone loss in celiac disease is related to secondary hyperparathyroidism. *J Bone Miner Res* 1999; 14: 652–7.
- 47 Pazianas M, Butcher GP, Subhani JM, *et al.* Calcium absorption and bone mineral density in celiacs after long term treatment with gluten-free diet and adequate calcium intake. *Osteoporos Int* 2005; 16: 56–63.
- 48 Taranta A, Fortunati D, Longo M, *et al.* Imbalance of osteoclastogenesis-regulating factors in patients with celiac disease. *J Bone Miner Res* 2004; 19: 1112–21.
- 49 Walters JR. Bone mineral density in coeliac disease. *Gut* 1994; 35: 150–1.
- 50 Corazza GR, Di Stefano M, Maurino E, *et al.* Bones in celiac disease: diagnosis and treatment. *Best Pract Res Clin Gastroenterol* 2005; 19: 453–65.
- 51 Cranney A, Rostom A, Sy R, *et al.* Consequences of testing for celiac disease. *Gastroenterology* 2005; 128: S109–20.
- 52 Valdimarsson T, Toss G, Lofman O, *et al.* Three years' follow-up of bone density in adult coeliac disease: significance of secondary hyperparathyroidism. *Scand J Gastroenterol* 2000; 35: 274–80.
- 53 Sategna-Guidetti C, Grosso SB, Grosso S, *et al.* The effects of 1-year gluten withdrawal on bone mass, bone metabolism and nutritional status in newly-diagnosed adult coeliac disease patients. *Aliment Pharmacol Ther* 2000; 14: 35–43.
- 54 Ciacci C, Maurelli L, Klain M, *et al.* Effects of dietary treatment on bone mineral density in adults with celiac disease: factors predicting response. *Am J Gastroenterol* 1997; 92: 992–6.
- 55 Mustalahti K, Collin P, Sievanen H, *et al.* Osteopenia in patients with clinically silent coeliac disease warrants screening. *Lancet* 1999; 354: 744–5.
- 56 McFarlane XA, Bhalla AK, Robertson DA. Effect of a gluten free diet on osteopenia in adults with newly diagnosed coeliac disease. *Gut* 1996; 39: 180–4.
- 57 Mautalen C, Gonzalez D, Mazure R, *et al.* Effect of treatment on bone mass,

- mineral metabolism, and body composition in untreated celiac disease patients. *Am J Gastroenterol* 1997; 92: 313–8.
- 58 Mora S, Barera G, Ricotti A, *et al.* Reversal of low bone density with a gluten-free diet in children and adolescents with celiac disease. *Am J Clin Nutr* 1998; 67: 477–81.
- 59 Kempainen T, Kroger H, Janatuinen E, *et al.* Bone recovery after a gluten-free diet: a 5-year follow-up study. *Bone* 1999; 25: 355–60.
- 60 Barera G, Beccio S, Proverbio MC, *et al.* Longitudinal changes in bone metabolism and bone mineral content in children with celiac disease during consumption of a gluten-free diet. *Am J Clin Nutr* 2004; 79: 148–54.
- 61 Scott BB, Losowsky MS. Depressed cell-mediated immunity in coeliac disease. *Gut* 1976; 17: 900–5.
- 62 O'Donoghue DP, Lancaster-Smith M, Lavinier P, *et al.* T cell depletion in untreated adult coeliac disease. *Gut* 1976; 17: 328–31.
- 63 Di Sabatino A, Bertrandi E, Casadei Maldini M, *et al.* Phenotyping of peripheral blood lymphocytes in adult coeliac disease. *Immunology* 1998; 95: 572–6.
- 64 Di Sabatino A, D'Alo S, Millimaggi D, *et al.* Apoptosis and peripheral blood lymphocyte depletion in coeliac disease. *Immunology* 2001; 103: 435–40.
- 65 Vaknin A, Eliakim R, Ackerman Z, *et al.* Neurological abnormalities associated with celiac disease. *J Neurol* 2004; 251: 1393–7.
- 66 Kalaydjian AE, Eaton W, Cascella N, *et al.* The gluten connection: the association between schizophrenia and celiac disease. *Acta Psychiatr Scand* 2006; 113: 82–90.
- 67 Ashkenazi A, Krasilovsky D, Levin S, *et al.* Immunologic reaction of psychotic patients to fractions of gluten. *Am J Psychiatry* 1979; 136: 1306–9.
- 68 Pynnonen PA, Isometsa ET, Verkasalo MA, *et al.* Gluten-free diet may alleviate depressive and behavioural symptoms in adolescents with coeliac disease: a prospective follow-up case-series study. *BMC Psychiatry* 2005; 5: 14.
- 69 Hernanz A, Polanco I. Plasma precursor amino acids of central nervous system monoamines in children with coeliac disease. *Gut* 1991; 32: 1478–81.
- 70 Sollid LM. Coeliac disease: dissecting a complex inflammatory disorder. *Nat Rev Immunol* 2002; 2: 647–55.
- 71 Brown RR. Metabolism and biology of tryptophan. In: Fillippini G, Costa C, Bertazzo A, eds. *Recent Advances in Tryptophan Research: Tryptophan and Serotonergic Pathways*. New York: Plenum, 1996: 15–25.
- 72 Mention JJ, Ben Ahmed M, Begue B, *et al.* Interleukin 15: a key to disrupted intraepithelial lymphocyte homeostasis and lymphomagenesis in celiac disease. *Gastroenterology* 2003; 125: 730–45.
- 73 Cellier C, Delabesse E, Helmer C, *et al.* Refractory sprue, coeliac disease, and enteropathy-associated T-cell lymphoma. French Coeliac Disease Study Group. *Lancet* 2000; 356: 203–8.
- 74 Krauss N, Schuppan D. Monitoring non-responsive patients who have celiac disease. *Gastrointest Endosc Clin N Am* 2006; 16: 317–27.
- 75 Juby LD, Dixon MF, Axon AT. Abnormal intestinal permeability and jejunal morphology. *J Clin Pathol* 1987; 40: 714–8.
- 76 Drago S, El Asmar R, Di Pierro M, *et al.* Gliadin, zonulin and gut permeability: effects on celiac and non-celiac intestinal mucosa and intestinal cell lines. *Scand J Gastroenterol* 2006; 41: 408–19.
- 77 Fasano A. Intestinal zonulin: open sesame! *Gut* 2001; 49: 159–62.
- 78 Fasano A. Systemic autoimmune disorders in celiac disease. *Curr Opin Gastroenterol* 2006; 22: 674–9.
- 79 Arrieta MC, Bistriz L, Meddings JB. Alterations in intestinal permeability. *Gut* 2006; 55: 1512–20.
- 80 Volta U, Granito A, De Franceschi L, *et al.* Anti tissue transglutaminase antibodies as predictors of silent coeliac disease in patients with hypertransaminasaemia of unknown origin. *Dig Liver Dis* 2001; 33: 420–5.
- 81 Pelaez-Luna M, Schmulson M, Robles-Diaz G. Intestinal involvement is not sufficient to explain hypertransaminasaemia in celiac disease? *Med Hypotheses* 2005; 65: 937–41.
- 82 Novacek G, Miehsler W, Wrba F, *et al.* Prevalence and clinical importance of hypertransaminasaemia in coeliac disease. *Eur J Gastroenterol Hepatol* 1999; 11: 283–8.
- 83 Ackerman Z, Eliashiv S, Reches A, *et al.* Neurological manifestations in celiac disease and vitamin E deficiency. *J Clin Gastroenterol* 1989; 11: 603–5.
- 84 Shepherd SJ. *Dietary Management of Gastrointestinal Disease: Gluten in Coeliac Disease and FODMAPs in IBS*. PhD Thesis, Clayton: Monash University, 2008.
- 85 Saibeni S, Lecchi A, Meucci G, *et al.* Prevalence of hyperhomocysteinemia in adult gluten-sensitive enteropathy at diagnosis: role of B12, folate, and genetics. *Clin Gastroenterol Hepatol* 2005; 3: 574–80.
- 86 Kempainen T, Kroger H, Janatuinen E, *et al.* Osteoporosis in adult patients with celiac disease. *Bone* 1999; 24: 249–55.
- 87 McFarlane XA, Bhalla AK, Reeves DE, *et al.* Osteoporosis in treated adult coeliac disease. *Gut* 1995; 36: 710–4.
- 88 Hinks LJ, Inwards KD, Lloyd B, *et al.* Body content of selenium in coeliac disease. *Br Med J (Clin Res Ed)* 1984; 288: 1862–3.
- 89 Chanoine JP, Neve J, Wu S, *et al.* Selenium decreases thyroglobulin concentrations but does not affect the increased thyroxine-to-triiodothyronine ratio in children with congenital hypothyroidism. *J Clin Endocrinol Metab* 2001; 86: 1160–3.
- 90 Mauro A, Orsi L, Mortara P, *et al.* Cerebellar syndrome in adult celiac disease with vitamin E deficiency. *Acta Neurol Scand* 1991; 84: 167–70.
- 91 Murray JA, Moore SB, Van Dyke CT, *et al.* HLA DQ gene dosage and risk and severity of celiac disease. *Clin Gastroenterol Hepatol* 2007; 5: 1406–12.
- 92 Ciacci C, Cirillo M, Cavallaro R, *et al.* Long-term follow-up of celiac adults on gluten-free diet: prevalence and correlates of intestinal damage. *Digestion* 2002; 66: 178–85.
- 93 Fine KD, Meyer RL, Lee EL. The prevalence and causes of chronic diarrhea in patients with celiac sprue treated with a gluten-free diet. *Gastroenterology* 1997; 112: 1830–8.
- 94 Kluge F, Koch HK, Grosse-Wilde H, *et al.* Follow-up of treated adult celiac disease: clinical and morphological studies. *Hepatogastroenterology* 1982; 29: 17–23.
- 95 Green PH, Cellier C. Celiac disease. *N Engl J Med* 2007; 357: 1731–43.
- 96 Dickey W, Hughes DF, McMillan SA. Disappearance of endomysial antibodies in treated celiac disease does not indicate histological recovery. *Am J Gastroenterol* 2000; 95: 712–4.
- 97 Kaukinen K, Sulkanen S, Maki M, *et al.* IgA-class transglutaminase antibodies in evaluating the efficacy of gluten-free diet in coeliac disease. *Eur J Gastroenterol Hepatol* 2002; 14: 311–5.
- 98 Vahedi K, Mascart F, Mary JY, *et al.* Reliability of antitransglutaminase antibodies as predictors of gluten-free diet

- compliance in adult celiac disease. *Am J Gastroenterol* 2003; **98**: 1079–87.
- 99 Leffler DA, Edwards George JB, Dennis M, *et al.* A prospective comparative study of five measures of gluten-free diet adherence in adults with coeliac disease. *Aliment Pharmacol Ther* 2007; **26**: 1227–35.
- 100 Tursi A, Brandimarte G, Giorgetti GM. Lack of usefulness of anti-transglutaminase antibodies in assessing histologic recovery after gluten-free diet in celiac disease. *J Clin Gastroenterol* 2003; **37**: 387–91.
- 101 Valentini RA, Andreani ML, Corazza GR, *et al.* IgA endomysium antibody: a valuable tool in the screening of coeliac disease but not its follow-up. *Ital J Gastroenterol* 1994; **26**: 279–82.
- 102 Midhagen G, Aberg AK, Olcen P, *et al.* Antibody levels in adult patients with coeliac disease during gluten-free diet: a rapid initial decrease of clinical importance. *J Intern Med* 2004; **256**: 519–24.
- 103 Sategna-Guidetti C, Pulitano R, Grosso S, *et al.* Serum IgA antiendomysium antibody titers as a marker of intestinal involvement and diet compliance in adult celiac sprue. *J Clin Gastroenterol* 1993; **17**: 123–7.
- 104 Maki M, Collin P. Coeliac disease. *Lancet* 1997; **349**: 1755–9.
- 105 Ansaldi N, Tavassoli K, Fausone D, *et al.* Clinico-histological behavior of celiac patients after gluten load following the definitive diagnosis. *Pediatr Med Chir* 1988; **10**: 3–6.
- 106 Matysiak-Budnik T, Malamut G, de Serre NP, *et al.* Long-term follow-up of 61 coeliac patients diagnosed in childhood: evolution toward latency is possible on a normal diet. *Gut* 2007; **56**: 1379–86.
- 107 Tursi A, Brandimarte G, Giorgetti GM, *et al.* Endoscopic and histological findings in the duodenum of adults with celiac disease before and after changing to a gluten-free diet: a 2-year prospective study. *Endoscopy* 2006; **38**: 702–7.
- 108 Wahab PJ, Meijer JW, Mulder CJ. Histologic follow-up of people with celiac disease on a gluten-free diet: slow and incomplete recovery. *Am J Clin Pathol* 2002; **118**: 459–63.
- 109 Walker-Smith JA, Guandalini S, Schmitz J, *et al.* Revised criteria for diagnosis of coeliac disease. *Arch Dis Child* 1990; **65**: 909–11.
- 110 Mee AS, Burke M, Vallon AG, *et al.* Small bowel biopsy for malabsorption: comparison of the diagnostic adequacy of endoscopic forceps and capsule biopsy specimens. *Br Med J (Clin Res Ed)* 1985; **291**: 769–72.
- 111 Pais WP, Duerksen DR, Pettigrew NM, *et al.* How many duodenal biopsy specimens are required to make a diagnosis of celiac disease? *Gastrointest Endosc* 2008; **167**: 1082–7.
- 112 Hopper AD, Cross SS, Sanders DS. Patchy villous atrophy in adult patients with suspected gluten-sensitive enteropathy: is a multiple duodenal biopsy strategy appropriate? *Endoscopy* 2008; **40**: 219–24.
- 113 Banerjee R, Shekharan A, Ramji C, *et al.* Role of magnification endoscopy in the diagnosis and evaluation of suspected celiac disease: correlation with histology. *Indian J Gastroenterol* 2007; **26**: 67–9.
- 114 Lo A, Guelrud M, Essendorf H, *et al.* Classification of villous atrophy with enhanced magnification endoscopy in patients with celiac disease and tropical sprue. *Gastrointest Endosc* 2007; **66**: 377–82.
- 115 Zambelli A, Villanacci V, Buscarini E, *et al.* Confocal laser endomicroscopy in celiac disease: description of findings in two cases. *Endoscopy* 2007; **39**: 1018–20.
- 116 Trovato C, Sonzogni A, Ravizza D, *et al.* Celiac disease: *in vivo* diagnosis by confocal endomicroscopy. *Gastrointest Endosc* 2007; **65**: 1096–9.
- 117 Murray JA, Rubio-Tapia A, Van Dyke CT, *et al.* Mucosal atrophy in celiac disease: extent of involvement, correlation with clinical presentation, and response to treatment. *Clin Gastroenterol Hepatol* 2008; **6**: 186–93; quiz 25.
- 118 Lee SK, Lo W, Memeo L, *et al.* Duodenal histology in patients with celiac disease after treatment with a gluten-free diet. *Gastrointest Endosc* 2003; **57**: 187–91.
- 119 Bardella MT, Velio P, Cesana BM, *et al.* Coeliac disease: a histological follow-up study. *Histopathology* 2007; **50**: 465–71.
- 120 Wahab PJ, Crusius JB, Meijer JW, *et al.* Gluten challenge in borderline gluten-sensitive enteropathy. *Am J Gastroenterol* 2001; **96**: 1464–9.
- 121 Tooley KL, Saxon BR, Webster J, *et al.* A novel non-invasive biomarker for assessment of small intestinal mucositis in children with cancer undergoing chemotherapy. *Cancer Biol Ther* 2006; **5**: 1275–81.
- 122 Clarke JM, Pelton NC, Bajka BH, *et al.* Use of the ¹³C-sucrose breath test to assess chemotherapy-induced small intestinal mucositis in the rat. *Cancer Biol Ther* 2006; **5**: 34–8.
- 123 Papadia C, Sherwood RA, Kalantzis C, *et al.* Plasma citrulline concentration: a reliable marker of small bowel absorptive capacity independent of intestinal inflammation. *Am J Gastroenterol* 2007; **102**: 1474–82.
- 124 Vilela EG, de Abreu Ferrari Mde L, de Gama Torres HO, *et al.* Intestinal permeability and antigliadin antibody test for monitoring adult patients with celiac disease. *Dig Dis Sci* 2007; **52**: 1304–9.
- 125 Duerksen DR, Wilhelm-Boyles C, Parry DM. Intestinal permeability in long-term follow-up of patients with celiac disease on a gluten-free diet. *Dig Dis Sci* 2005; **50**: 785–90.
- 126 Cummins AG, Penttila IA, Labrooy JT, *et al.* Recovery of the small intestine in coeliac disease on a gluten-free diet: changes in intestinal permeability, small bowel morphology and T-cell activity. *J Gastroenterol Hepatol* 1991; **6**: 53–7.
- 127 Hallert C, Grant C, Grehn S, *et al.* Evidence of poor vitamin status in coeliac patients on a gluten-free diet for 10 years. *Aliment Pharmacol Ther* 2002; **16**: 1333–9.
- 128 Bardella MT, Molteni N, Prampolini L, *et al.* Need for follow up in coeliac disease. *Arch Dis Child* 1994; **70**: 211–3.
- 129 Lamontagne P, West GE, Galibois I. Quebecers with celiac disease: analysis of dietary problems. *Can J Diet Pract Res* 2001; **62**: 175–81.
- 130 Hogberg L, Grodzinsky E, Stenhammar L. Better dietary compliance in patients with coeliac disease diagnosed in early childhood. *Scand J Gastroenterol* 2003; **38**: 751–4.
- 131 Ljungman G, Myrdal U. Compliance in teenagers with coeliac disease—a Swedish follow-up study. *Acta Paediatr* 1993; **82**: 235–8.
- 132 Bebb JR, Lawson A, Knight T, *et al.* Long-term follow-up of coeliac disease—what do coeliac patients want? *Aliment Pharmacol Ther* 2006; **23**: 827–31.
- 133 Pietzak MM. Follow-up of patients with celiac disease: achieving compliance with treatment. *Gastroenterology* 2005; **128**: S135–41.
- 134 Holmes GK. Neurological and psychiatric complications in coeliac disease. In: Gobbi G, Andermann F, Naccarato S, *et al.*, eds. *Epilepsy and Other Neurological Disorders in Coeliac Disease*. London: John Libbey, 1997: 251–64.
- 135 Ciacci C, Iavarone A, Siniscalchi M, *et al.* Psychological dimensions of celiac disease: toward an integrated approach. *Dig Dis Sci* 2002; **47**: 2082–7.

- 136 Bode S, Gudmand-Hoyer E. Symptoms and haematologic features in consecutive adult coeliac patients. *Scand J Gastroenterol* 1996; 31: 54–60.
- 137 Marshall D, Johnell O, Wedel H. Meta-analysis of how well measures of bone mineral density predict occurrence of osteoporotic fractures. *BMJ* 1996; 312: 1254–9.
- 138 van Hemert AM, Vandenbroucke JP, Birkenhager JC, *et al.* Prediction of osteoporotic fractures in the general population by a fracture risk score. A 9-year follow-up among middle-aged women. *Am J Epidemiol* 1990; 132: 123–35.
- 139 van Heel DA, West J. Recent advances in coeliac disease. *Gut* 2006; 55: 1037–46.
- 140 Lewis NR, Scott BB. Should patients with coeliac disease have their bone mineral density measured? *Eur J Gastroenterol Hepatol* 2005; 17: 1065–70.
- 141 Logan R, West J. Risk of fracture in coeliac disease. *Gut* 2003; 52: 1532.
- 142 Catassi C, Fabiani E, Iacono G, *et al.* A prospective, double-blind, placebo-controlled trial to establish a safe gluten threshold for patients with celiac disease. *Am J Clin Nutr* 2007; 85: 160–6.
- 143 Akobeng AK, Thomas AG. Systematic review: tolerable amount of gluten for people with coeliac disease. *Aliment Pharmacol Ther* 2008; 27: 1044–52.
- 144 Garsed K, Scott BB. Can oats be taken in a gluten-free diet? A systematic review. *Scand J Gastroenterol* 2007; 42: 171–8.
- 145 Rashid M, Butzner D, Burrows V, *et al.* Consumption of pure oats by individuals with coeliac disease: a position statement by the Canadian Celiac Association. *Can J Gastroenterol* 2007; 21: 649–51.
- 146 Ellis HJ, Ciclitira PJ. Should coeliac sufferers be allowed their oats? *Eur J Gastroenterol Hepatol* 2008; 20: 492–3.
- 147 Silano M, Dessi M, De Vincenzi M, *et al.* *In vitro* tests indicate that certain varieties of oats may be harmful to patients with coeliac disease. *J Gastroenterol Hepatol* 2007; 22: 528–31.
- 148 Hernando A, Mujico JR, Mena MC, *et al.* Measurement of wheat gluten and barley hordeins in contaminated oats from Europe, the United States and Canada by Sandwich R5 ELISA. *Eur J Gastroenterol Hepatol* 2008; 20: 545–54.
- 149 Barrett JS, Gibson PR. Clinical ramifications of malabsorption of fructose and other short-chain carbohydrates. *Pract Gastroenterol* 2007; 31: 51–65.
- 150 Patey-Mariaud De Serre N, Cellier C, Jabri B, *et al.* Distinction between coeliac disease and refractory sprue: a simple immunohistochemical method. *Histopathology* 2000; 37: 70–7.
- 151 Cellier C, Patey N, Mauvieux L, *et al.* Abnormal intestinal intraepithelial lymphocytes in refractory sprue. *Gastroenterology* 1998; 114: 471–81.
- 152 Daum S, Cellier C, Mulder CJ. Refractory coeliac disease. *Best Pract Res Clin Gastroenterol* 2005; 19: 413–24.
- 153 Tomei E, Diacinti D, Marini M, *et al.* Abdominal CT findings may suggest coeliac disease. *Dig Liver Dis* 2005; 37: 402–6.
- 154 Hadithi M, Mallant M, Oudejans J, *et al.* 18F-FDG PET versus CT for the detection of enteropathy-associated T-cell lymphoma in refractory celiac disease. *J Nucl Med* 2006; 47: 1622–7.
- 155 Hadithi M, Al-toma A, Oudejans J, *et al.* The value of double-balloon enteroscopy in patients with refractory celiac disease. *Am J Gastroenterol* 2007; 102: 987–96.
- 156 Vanderpump MJ, Tunbridge WMG. The epidemiology of thyroid disease. In: Braverman LE, Utiger RD, eds. *The Thyroid: A Fundamental and Clinical Text*. Philadelphia, PA: Lippincott-Raven, 1996: 474.
- 157 Bottaro G, Cataldo F, Rotolo N, *et al.* The clinical pattern of subclinical/silent celiac disease: an analysis on 1026 consecutive cases. *Am J Gastroenterol* 1999; 94: 691–6.
- 158 Sategna-Guidetti C, Volta U, Ciacci C, *et al.* Prevalence of thyroid disorders in untreated adult celiac disease patients and effect of gluten withdrawal: an Italian multicenter study. *Am J Gastroenterol* 2001; 96: 751–7.
- 159 Hakonen M, Luotola K, Salmi J, *et al.* Clinical and subclinical autoimmune thyroid disease in adult celiac disease. *Dig Dis Sci* 2001; 46: 2631–5.
- 160 Counsell CE, Taha A, Ruddell WS. Coeliac disease and autoimmune thyroid disease. *Gut* 1994; 35: 844–6.
- 161 Ansaldi N, Palmas T, Corrias A, *et al.* Autoimmune thyroid disease and celiac disease in children. *J Pediatr Gastroenterol Nutr* 2003; 37: 63–6.
- 162 Kowalska E, Wasowska-Krolukowska K, Toporowska-Kowalska E. Estimation of antithyroid antibodies occurrence in children with coeliac disease. *Med Sci Monit* 2000; 6: 719–21.
- 163 Mankai A, Chadli-Chaieb M, Saad F, *et al.* Screening for celiac disease in Tunisian patients with Graves' disease using anti-endomysium and anti-tissue transglutaminase antibodies. *Gastroenterol Clin Biol* 2006; 30: 961–4.
- 164 Cuoco L, Certo M, Jorizzo RA, *et al.* Prevalence and early diagnosis of coeliac disease in autoimmune thyroid disorders. *Ital J Gastroenterol Hepatol* 1999; 31: 283–7.
- 165 Spadaccino AC, Basso D, Chiarelli S, *et al.* Celiac disease in North Italian patients with autoimmune thyroid diseases. *Autoimmunity* 2008; 41: 116–21.
- 166 Guliter S, Yakaryilmaz F, Ozkurt Z, *et al.* Prevalence of coeliac disease in patients with autoimmune thyroiditis in a Turkish population. *World J Gastroenterol* 2007; 13: 1599–601.
- 167 Toscano V, Conti FG, Anastasi E, *et al.* Importance of gluten in the induction of endocrine autoantibodies and organ dysfunction in adolescent celiac patients. *Am J Gastroenterol* 2000; 95: 1742–8.
- 168 Fracchia M, Galatola G, Corradi F, *et al.* Coeliac disease associated with Sjogren's syndrome, renal tubular acidosis, primary biliary cirrhosis and autoimmune hyperthyroidism. *Dig Liver Dis* 2004; 36: 489–91.
- 169 Pais WP, Duerksen DR, Pettigrew NM, *et al.* How many duodenal biopsy specimens are required to make a diagnosis of celiac disease? *Gastrointest Endosc* 2008; 67: 1082–7.
- 170 Collin P, Kaukinen K, Valimaki M, *et al.* Endocrinological disorders and celiac disease. *Endocr Rev* 2002; 23: 464–83.
- 171 de Freitas IN, Sipahi AM, Damiao AO, *et al.* Celiac disease in Brazilian adults. *J Clin Gastroenterol* 2002; 34: 430–4.
- 172 Goh C, Banerjee K. Prevalence of coeliac disease in children and adolescents with type 1 diabetes mellitus in a clinic based population. *Postgrad Med J* 2007; 83: 132–6.
- 173 Aktay AN, Lee PC, Kumar V, *et al.* The prevalence and clinical characteristics of celiac disease in juvenile diabetes in Wisconsin. *J Pediatr Gastroenterol Nutr* 2001; 33: 462–5.
- 174 Cronin CC, Feighery A, Ferriss JB, *et al.* High prevalence of celiac disease among patients with insulin-dependent (type I) diabetes mellitus. *Am J Gastroenterol* 1997; 92: 2210–2.
- 175 Not T, Tommasini A, Tonini G, *et al.* Undiagnosed coeliac disease and risk of autoimmune disorders in subjects with Type I diabetes mellitus. *Diabetologia* 2001; 44: 151–5.
- 176 Ludvigsson JF, Ludvigsson J, Ekblom A, *et al.* Celiac disease and risk of subsequent type 1 diabetes: a general population cohort study of children and

- adolescents. *Diabetes Care* 2006; **29**: 2483–8.
- 177 Anonymous. Prevalence of diabetes among Hispanics – selected areas, 1998–2002. *Morb Mortal Wkly Rep* 2004; **53**: 941–4.
- 178 Sanchez JC, Cabrera-Rode E, Sorell L, *et al.* Celiac disease associated antibodies in persons with latent autoimmune diabetes of adult and type 2 diabetes. *Autoimmunity* 2007; **40**: 103–7.
- 179 Harris MI, Robbins DC. Prevalence of adult-onset IDDM in the U.S. population. *Diabetes Care* 1994; **17**: 1337–40.
- 180 Landin-Olsson M, Nilsson KO, Lernmark A, *et al.* Islet cell antibodies and fasting C-peptide predict insulin requirement at diagnosis of diabetes mellitus. *Diabetologia* 1990; **33**: 561–8.
- 181 Niskanen LK, Tuomi T, Karjalainen J, *et al.* GAD antibodies in NIDDM. Ten-year follow-up from the diagnosis. *Diabetes Care* 1995; **18**: 1557–65.
- 182 Volta U, De Franceschi L, Molinaro N, *et al.* Frequency and significance of anti-gliadin and anti-endomysial antibodies in autoimmune hepatitis. *Dig Dis Sci* 1998; **43**: 2190–5.
- 183 Villalta D, Girolami D, Bidoli E, *et al.* High prevalence of celiac disease in autoimmune hepatitis detected by anti-tissue transglutaminase autoantibodies. *J Clin Lab Anal* 2005; **19**: 6–10.
- 184 Caprai S, Vajro P, Ventura A, *et al.* Autoimmune liver disease associated with celiac disease in childhood: a multicenter study. *Clin Gastroenterol Hepatol* 2008; **6**: 803–6.
- 185 Sjoberg K, Lindgren S, Eriksson S. Frequent occurrence of non-specific gliadin antibodies in chronic liver disease. Endomysial but not gliadin antibodies predict coeliac disease in patients with chronic liver disease. *Scand J Gastroenterol* 1997; **32**: 1162–7.
- 186 Kabasakal Y, Kitapcioglu G, Turk T, *et al.* The prevalence of Sjogren's syndrome in adult women. *Scand J Rheumatol* 2006; **35**: 379–83.
- 187 Thomas E, Hay EM, Hajeer A, *et al.* Sjogren's syndrome: a community-based study of prevalence and impact. *Br J Rheumatol* 1998; **37**: 1069–76.
- 188 Sanchez-Guerrero J, Perez-Dosal MR, Cardenas-Velazquez F, *et al.* Prevalence of Sjogren's syndrome in ambulatory patients according to the American-European Consensus Group criteria. *Rheumatology (Oxford)* 2005; **44**: 235–40.
- 189 Szodoray P, Barta Z, Lakos G, *et al.* Coeliac disease in Sjogren's syndrome – a study of 111 Hungarian patients. *Rheumatol Int* 2004; **24**: 278–82.
- 190 Ludvigsson JF, Montgomery SM, Olen O, *et al.* Coeliac disease and risk of renal disease—a general population cohort study. *Nephrol Dial Transplant* 2006; **21**: 1809–15.
- 191 Matteoni CA, Goldblum JR, Wang N, *et al.* Celiac disease is highly prevalent in lymphocytic colitis. *J Clin Gastroenterol* 2001; **32**: 225–7.
- 192 Fine KD, Do K, Schulte K, *et al.* High prevalence of celiac sprue-like HLA-DQ genes and enteropathy in patients with the microscopic colitis syndrome. *Am J Gastroenterol* 2000; **95**: 1974–82.
- 193 McCashland TM, Donovan JP, Strobach RS, *et al.* Collagenous enterocolitis: a manifestation of gluten-sensitive enteropathy. *J Clin Gastroenterol* 1992; **15**: 45–51.
- 194 Willis AC, Vince FP. The prevalence of Addison's disease in Coventry, UK. *Postgrad Med J* 1997; **73**: 286–8.
- 195 Biagi F, Campanella J, Soriani A, *et al.* Prevalence of coeliac disease in Italian patients affected by Addison's disease. *Scand J Gastroenterol* 2006; **41**: 302–5.
- 196 Elfstrom P, Montgomery SM, Kampe O, *et al.* Risk of primary adrenal insufficiency in patients with celiac disease. *J Clin Endocrinol Metab* 2007; **92**: 3595–8.
- 197 Myhre AG, Undlien DE, Lovas K, *et al.* Autoimmune adrenocortical failure in Norway autoantibodies and human leukocyte antigen class II associations related to clinical features. *J Clin Endocrinol Metab* 2002; **87**: 618–23.
- 198 O'Leary C, Walsh CH, Wieneke P, *et al.* Coeliac disease and autoimmune Addison's disease: a clinical pitfall. *QJM* 2002; **95**: 79–82.
- 199 Bernatsky S, Joseph L, Pineau CA, *et al.* A population-based assessment of systemic lupus erythematosus incidence and prevalence – results and implications of using administrative data for epidemiological studies. *Rheumatology (Oxford)* 2007; **46**: 1814–8.
- 200 Benucci M, Del Rosso A, Li Gobbi F, *et al.* Systemic lupus erythematosus (SLE) in Italy: an Italian prevalence study based on a two-step strategy in an area of Florence (Scandicci-Le Signe). *Med Sci Monit* 2005; **11**: CR420–5.
- 201 Molina MJ, Mayor AM, Franco AE, *et al.* Prevalence of systemic lupus erythematosus and associated comorbidities in Puerto Rico. *J Clin Rheumatol* 2007; **13**: 202–4.
- 202 Mirza N, Bonilla E, Phillips PE. Celiac disease in a patient with systemic lupus erythematosus: a case report and review of literature. *Clin Rheumatol* 2007; **26**: 827–8.
- 203 Madani S, Shapiro J. Alopecia areata update. *J Am Acad Dermatol* 2000; **42**: 549–66; quiz 67–70.
- 204 Kundrotas LW, Clement DJ. Serum alanine aminotransferase (ALT) elevation in asymptomatic US Air Force basic trainee blood donors. *Dig Dis Sci* 1993; **38**: 2145–50.
- 205 Ioannou GN, Boyko EJ, Lee SP. The prevalence and predictors of elevated serum aminotransferase activity in the United States in 1999–2002. *Am J Gastroenterol* 2006; **101**: 76–82.
- 206 Carroccio A, Di Prima L, Scalici C, *et al.* Unexplained elevated serum pancreatic enzymes: a reason to suspect celiac disease. *Clin Gastroenterol Hepatol* 2006; **4**: 455–9.
- 207 Farre C, Esteve M, Curcoy A, *et al.* Hypertransaminasemia in pediatric celiac disease patients and its prevalence as a diagnostic clue. *Am J Gastroenterol* 2002; **97**: 3176–81.
- 208 Hagander B, Berg NO, Brandt L, *et al.* Hepatic injury in adult coeliac disease. *Lancet* 1977; **2**: 270–2.
- 209 Bardella MT, Fraquelli M, Quatrini M, *et al.* Prevalence of hypertransaminasemia in adult celiac patients and effect of gluten-free diet. *Hepatology* 1995; **22**: 833–6.
- 210 Bardella MT, Quatrini M, Zuin M, *et al.* Screening patients with celiac disease for primary biliary cirrhosis and vice versa. *Am J Gastroenterol* 1997; **92**: 1524–6.
- 211 Volta U, De Franceschi L, Lari F, *et al.* Coeliac disease hidden by cryptogenic hypertransaminasaemia. *Lancet* 1998; **352**: 26–9.
- 212 Bardella MT, Vecchi M, Conte D, *et al.* Chronic unexplained hypertransaminasemia may be caused by occult celiac disease. *Hepatology* 1999; **29**: 654–7.
- 213 Myszor M, James OF. The epidemiology of primary biliary cirrhosis in north-east England: an increasingly common disease? *Q J Med* 1990; **75**: 377–85.
- 214 James OF, Bhopal R, Howel D, *et al.* Primary biliary cirrhosis once rare, now common in the United Kingdom? *Hepatology* 1999; **30**: 390–4.
- 215 Floreani A, Betterle C, Baragiotta A, *et al.* Prevalence of coeliac disease in primary biliary cirrhosis and of anti-mitochondrial antibodies in adult coeliac

- disease patients in Italy. *Dig Liver Dis* 2002; 34: 258–61.
- 216 Chatzicostas C, Roussomoustakaki M, Drygiannakis D, *et al.* Primary biliary cirrhosis and autoimmune cholangitis are not associated with coeliac disease in Crete. *BMC Gastroenterol* 2002; 2: 5.
- 217 Niveloni S, Dezi R, Pedreira S, *et al.* Gluten sensitivity in patients with primary biliary cirrhosis. *Am J Gastroenterol* 1998; 93: 404–8.
- 218 Rubio-Tapia A, Murray JA. The liver in coeliac disease. *Hepatology* 2007; 46: 1650–8.
- 219 Volta U, Rodrigo L, Granito A, *et al.* Celiac disease in autoimmune cholestatic liver disorders. *Am J Gastroenterol* 2002; 97: 2609–13.
- 220 Ludvigsson JF, Elfstrom P, Broome U, *et al.* Celiac disease and risk of liver disease: a general population-based study. *Clin Gastroenterol Hepatol* 2007; 5: 63–9.
- 221 Habiore A, Lewartowska A, Orlowska J, *et al.* Association of coeliac disease with primary biliary cirrhosis in Poland. *Eur J Gastroenterol Hepatol* 2003; 15: 159–64.
- 222 Lindgren S, Sjoberg K, Eriksson S. Unsuspected coeliac disease in chronic 'cryptogenic' liver disease. *Scand J Gastroenterol* 1994; 29: 661–4.
- 223 Ojetti V, Fini L, Zileri Dal Verme L, *et al.* Acute cryptogenic liver failure in an untreated coeliac patient: a case report. *Eur J Gastroenterol Hepatol* 2005; 17: 1119–21.
- 224 Bambha K, Kim WR, Talwalkar J, *et al.* Incidence, clinical spectrum, and outcomes of primary sclerosing cholangitis in a United States community. *Gastroenterology* 2003; 125: 1364–9.
- 225 Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med* 2002; 346: 1221–31.
- 226 Lo Iacono O, Petta S, Venezia G, *et al.* Anti-tissue transglutaminase antibodies in patients with abnormal liver tests: is it always coeliac disease? *Am J Gastroenterol* 2005; 100: 2472–7.
- 227 Bardella MT, Valenti L, Pagliari C, *et al.* Searching for coeliac disease in patients with non-alcoholic fatty liver disease. *Dig Liver Dis* 2004; 36: 333–6.
- 228 Garg PK, Tandon RK. Survey on chronic pancreatitis in the Asia-Pacific region. *J Gastroenterol Hepatol* 2004; 19: 998–1004.
- 229 Rothenbacher D, Low M, Hardt PD, *et al.* Prevalence and determinants of exocrine pancreatic insufficiency among older adults: results of a population-based study. *Scand J Gastroenterol* 2005; 40: 697–704.
- 230 Patel RS, Johlin FC Jr, Murray JA. Celiac disease and recurrent pancreatitis. *Gastrointest Endosc* 1999; 50: 823–7.
- 231 Ludvigsson JF, Montgomery SM, Ekblom A. Risk of pancreatitis in 14,000 individuals with celiac disease. *Clin Gastroenterol Hepatol* 2007; 5: 1347–53.
- 232 Wills A, Hovell CJ. Neurological complications of enteric disease. *Gut* 1996; 39: 501–4.
- 233 Cooke WT, Smith WT. Neurological disorders associated with adult coeliac disease. *Brain* 1966; 89: 683–722.
- 234 Kepes JJ, Chou SM, Price LW Jr. Progressive multifocal leukoencephalopathy with 10-year survival in a patient with nontropical sprue. Report of a case with unusual light and electron microscopic features. *Neurology* 1975; 25: 1006–12.
- 235 Collin P, Pirttila T, Nurmikko T, *et al.* Celiac disease, brain atrophy, and dementia. *Neurology* 1991; 41: 372–5.
- 236 Gobbi G, Bouquet F, Greco L, *et al.* Coeliac disease, epilepsy, and cerebral calcifications. The Italian Working Group on Coeliac Disease and Epilepsy. *Lancet* 1992; 340: 439–43.
- 237 Hadjivassiliou M, Chattopadhyay AK, Davies-Jones GA, *et al.* Neuromuscular disorder as a presenting feature of coeliac disease. *J Neurol Neurosurg Psychiatry* 1997; 63: 770–5.
- 238 Kaplan JG, Pack D, Horoupian D, *et al.* Distal axonopathy associated with chronic gluten enteropathy: a treatable disorder. *Neurology* 1988; 38: 642–5.
- 239 Hadjivassiliou M, Grunewald RA, Chattopadhyay AK, *et al.* Clinical, radiological, neurophysiological, and neuropathological characteristics of gluten ataxia. *Lancet* 1998; 352: 1582–5.
- 240 Hu WT, Murray JA, Greenaway MC, *et al.* Cognitive impairment and coeliac disease. *Arch Neurol* 2006; 63: 1440–6.
- 241 Rigamonti A, Magi S, Venturini E, *et al.* Celiac disease presenting with motor neuropathy: effect of gluten free-diet. *Muscle Nerve* 2007; 35: 675–7.
- 242 Wong D, Dwinell M, Schulzer M, *et al.* Ataxia and the role of antigliadin antibodies. *Can J Neurol Sci* 2007; 34: 193–6.
- 243 Ranua J, Luoma K, Auvinen A, *et al.* Celiac disease-related antibodies in an epilepsy cohort and matched reference population. *Epilepsy Behav* 2005; 6: 388–92.
- 244 Boeing L, Murray V, Pelosi A, *et al.* Adolescent-onset psychosis: prevalence, needs and service provision. *Br J Psychiatry* 2007; 190: 18–26.
- 245 Kendler KS, Gallagher TJ, Abelson JM, *et al.* Lifetime prevalence, demographic risk factors, and diagnostic validity of nonaffective psychosis as assessed in a US community sample. The National Comorbidity Survey. *Arch Gen Psychiatry* 1996; 53: 1022–31.
- 246 Ludvigsson JF, Osby U, Ekblom A, *et al.* Coeliac disease and risk of schizophrenia and other psychosis: a general population cohort study. *Scand J Gastroenterol* 2007; 42: 179–85.
- 247 Eaton W, Mortensen PB, Agerbo E, *et al.* Coeliac disease and schizophrenia: population based case control study with linkage of Danish national registers. *BMJ* 2004; 328: 438–9.
- 248 West J, Logan RF, Hubbard RB, *et al.* Risk of schizophrenia in people with coeliac disease, ulcerative colitis and Crohn's disease: a general population-based study. *Aliment Pharmacol Ther* 2006; 23: 71–4.
- 249 Peleg R, Ben-Zion ZI, Peleg A, *et al.* "Bread madness" revisited: screening for specific celiac antibodies among schizophrenia patients. *Eur Psychiatry* 2004; 19: 311–4.
- 250 Compton WM, Conway KP, Stinson FS, *et al.* Changes in the prevalence of major depression and comorbid substance use disorders in the United States between 1991–1992 and 2001–2002. *Am J Psychiatry* 2006; 163: 2141–7.
- 251 Kessler RC, Chiu WT, Demler O, *et al.* Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005; 62: 617–27.
- 252 Anstey KJ, von Sanden C, Sargent-Cox K, *et al.* Prevalence and risk factors for depression in a longitudinal, population-based study including individuals in the community and residential care. *Am J Geriatr Psychiatry* 2007; 15: 497–505.
- 253 Blay SL, Andreoli SB, Fillenbaum GG, *et al.* Depression morbidity in later life: prevalence and correlates in a developing country. *Am J Geriatr Psychiatry* 2007; 15: 790–9.
- 254 Addolorato G, Capristo E, Ghittoni G, *et al.* Anxiety but not depression decreases in coeliac patients after one-year gluten-free diet: a longitudinal study. *Scand J Gastroenterol* 2001; 36: 502–6.

- 255 Ludvigsson JF, Reutfors J, Osby U, *et al.* Coeliac disease and risk of mood disorders – a general population-based cohort study. *J Affect Disord* 2007; **99**: 117–26.
- 256 Pynnönen PA, Isometsä ET, Verkasalo MA, *et al.* Untreated coeliac disease and development of mental disorders in children and adolescents. *Psychosomatics* 2002; **43**: 331–4.
- 257 Ciacci C, Iavarone A, Mazzacca G, *et al.* Depressive symptoms in adult coeliac disease. *Scand J Gastroenterol* 1998; **33**: 247–50.
- 258 Munk-Jørgensen P, Allgulander C, Dahl AA, *et al.* Prevalence of generalized anxiety disorder in general practice in Denmark, Finland, Norway, and Sweden. *Psychiatr Serv* 2006; **57**: 1738–44.
- 259 Somers JM, Goldner EM, Waraich P, *et al.* Prevalence and incidence studies of anxiety disorders: a systematic review of the literature. *Can J Psychiatry* 2006; **51**: 100–13.
- 260 Kroenke K, Spitzer RL, Williams JB, *et al.* Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Ann Intern Med* 2007; **146**: 317–25.
- 261 Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999; **281**: 537–44.
- 262 Feldman HA, Goldstein I, Hatzichristou DG, *et al.* Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Aging Study. *J Urol* 1994; **151**: 54–61.
- 263 Farthing MJ, Edwards CR, Rees LH, *et al.* Male gonadal function in coeliac disease: 1. Sexual dysfunction, infertility, and semen quality. *Gut* 1982; **23**: 608–14.
- 264 Ciacci C, De Rosa A, de Michele G, *et al.* Sexual behaviour in untreated and treated coeliac patients. *Eur J Gastroenterol Hepatol* 1998; **10**: 649–51.
- 265 McKinlay SM, Brambilla DJ, Posner JG. The normal menopause transition. *Maturitas* 1992; **14**: 103–15.
- 266 Munster K, Helm P, Schmidt L. Secondary amenorrhoea: prevalence and medical contact – a cross-sectional study from a Danish county. *Br J Obstet Gynaecol* 1992; **99**: 430–3.
- 267 Molteni N, Bardella MT, Bianchi PA. Obstetric and gynecological problems in women with untreated coeliac sprue. *J Clin Gastroenterol* 1990; **12**: 37–9.
- 268 Smecul E, Maurino E, Vazquez H, *et al.* Gynaecological and obstetric disorders in coeliac disease: frequent clinical onset during pregnancy or the puerperium. *Eur J Gastroenterol Hepatol* 1996; **8**: 63–89.
- 269 Pradhan M, Manisha, Singh R, *et al.* Coeliac disease as a rare cause of primary amenorrhoea: a case report. *J Reprod Med* 2007; **52**: 453–5.
- 270 Stephen EH, Chandra A. Declining estimates of infertility in the United States: 1982–2002. *Fertil Steril* 2006; **86**: 516–23.
- 271 Anderson MS, Hay WW. Intrauterine growth restriction and the small-for-gestational-age infant. In: Avery GB, Fletcher MA, MacDonald MG, eds. *Neonatology Pathophysiology and Management of the Newborn*. Philadelphia, PA: Lippincott Williams and Wilkins, 1999: 411.
- 272 Collin P, Vilska S, Heinonen PK, *et al.* Infertility and coeliac disease. *Gut* 1996; **39**: 382–4.
- 273 Meloni GF, Dessole S, Vargiu N, *et al.* The prevalence of coeliac disease in infertility. *Hum Reprod* 1999; **14**: 2759–61.
- 274 Rostami K, Steegers EA, Wong WY, *et al.* Coeliac disease and reproductive disorders: a neglected association. *Eur J Obstet Gynecol Reprod Biol* 2001; **96**: 146–9.
- 275 Martinelli P, Troncone R, Paparo F, *et al.* Coeliac disease and unfavourable outcome of pregnancy. *Gut* 2000; **46**: 332–5.
- 276 Ludvigsson JF, Montgomery SM, Ekbohm A. Coeliac disease and risk of adverse fetal outcome: a population-based cohort study. *Gastroenterology* 2005; **129**: 454–63.
- 277 Norgard B, Fonager K, Sorensen HT, *et al.* Birth outcomes of women with coeliac disease: a nationwide historical cohort study. *Am J Gastroenterol* 1999; **94**: 2435–40.
- 278 Sher KS, Mayberry JF. Female fertility, obstetric and gynaecological history in coeliac disease: a case control study. *Acta Paediatr Suppl* 1996; **412**: 76–7.
- 279 Ciacci C, Cirillo M, Auriemma G, *et al.* Coeliac disease and pregnancy outcome. *Am J Gastroenterol* 1996; **91**: 718–22.
- 280 Pellicano R, Astegiano M, Bruno M, *et al.* Women and coeliac disease: association with unexplained infertility. *Minerva Med* 2007; **98**: 217–9.
- 281 Tata LJ, Card TR, Logan RF, *et al.* Fertility and pregnancy-related events in women with coeliac disease: a population-based cohort study. *Gastroenterology* 2005; **128**: 849–55.
- 282 Jackson JE, Rosen M, McLean T, *et al.* Prevalence of coeliac disease in a cohort of women with unexplained infertility. *Fertil Steril* 2008; **89**: 1002–4.
- 283 Sheiner E, Peleg R, Levy A. Pregnancy outcome of patients with known coeliac disease. *Eur J Obstet Gynecol Reprod Biol* 2006; **129**: 41–5.
- 284 Kolho KL, Tiitinen A, Tulppala M, *et al.* Screening for coeliac disease in women with a history of recurrent miscarriage or infertility. *Br J Obstet Gynaecol* 1999; **106**: 171–3.
- 285 Smeets-Goevaers CG, Lesusink GL, Pappapoulos SE, *et al.* The prevalence of low bone mineral density in Dutch perimenopausal women: the Eindhoven perimenopausal osteoporosis study. *Osteoporos Int* 1998; **8**: 404–9.
- 286 Looker AC, Orwoll ES, Johnston CC Jr, *et al.* Prevalence of low femoral bone density in older U.S. adults from NHANES III. *J Bone Miner Res* 1997; **12**: 1761–8.
- 287 O'Leary C, Feighery C, Feighery A, *et al.* The prevalence of coeliac disease among female subjects having bone densitometry. *Ir J Med Sci* 2002; **171**: 145–7.
- 288 Fiore CE, Pennisi P, Ferro G, *et al.* Altered osteoprotegerin/RANKL ratio and low bone mineral density in coeliac patients on long-term treatment with gluten-free diet. *Horm Metab Res* 2006; **38**: 417–22.
- 289 Pistorius LR, Sweidan WH, Purdie DW, *et al.* Coeliac disease and bone mineral density in adult female patients. *Gut* 1995; **37**: 639–42.
- 290 Valdimarsson T, Toss G, Ross I, *et al.* Bone mineral density in coeliac disease. *Scand J Gastroenterol* 1994; **29**: 457–61.
- 291 Henry MJ, Pasco JA, Nicholson GC, *et al.* Prevalence of osteoporosis in Australian women: Geelong Osteoporosis Study. *J Clin Densitom* 2000; **3**: 261–8.
- 292 Nuti R, Martini G, Valenti R, *et al.* Prevalence of undiagnosed coeliac syndrome in osteoporotic women. *J Intern Med* 2001; **250**: 361–6.
- 293 Vasquez H, Mazure R, Gonzalez D, *et al.* Risk of fractures in coeliac disease patients: a cross-sectional, case-control study. *Am J Gastroenterol* 2000; **95**: 183–9.
- 294 Thomason K, West J, Logan RF, *et al.* Fracture experience of patients with coeliac disease: a population based survey. *Gut* 2003; **52**: 518–22.
- 295 Davie MW, Gaywood I, George E, *et al.* Excess non-spine fractures in women over 50 years with coeliac disease: a

- cross-sectional, questionnaire-based study. *Osteoporos Int* 2005; 16: 1150–5.
- 296 Olmos M, Antelo M, Vazquez H, *et al.* Systematic review and meta-analysis of observational studies on the prevalence of fractures in coeliac disease. *Dig Liver Dis* 2008; 40: 46–53.
- 297 Moreno ML, Vazquez H, Mazure R, *et al.* Stratification of bone fracture risk in patients with celiac disease. *Clin Gastroenterol Hepatol* 2004; 2: 127–34.
- 298 Ludvigsson JF, Michaelsson K, Ekblom A, *et al.* Coeliac disease and the risk of fractures – a general population-based cohort study. *Aliment Pharmacol Ther* 2007; 25: 273–85.
- 299 West J, Logan RF, Card TR, *et al.* Fracture risk in people with celiac disease: a population-based cohort study. *Gastroenterology* 2003; 125: 429–36.
- 300 Vestergaard P, Mosekilde L. Fracture risk in patients with celiac Disease, Crohn's disease, and ulcerative colitis: a nationwide follow-up study of 16,416 patients in Denmark. *Am J Epidemiol* 2002; 156: 1–10.
- 301 McCarthy CF, Fraser ID, Evans KT, *et al.* Lymphoreticular dysfunction in idiopathic steatorrhea. *Gut* 1966; 7: 140–8.
- 302 Corazza GR, Zoli G, Di Sabatino A, *et al.* A reassessment of splenic hypofunction in celiac disease. *Am J Gastroenterol* 1999; 94: 391–7.
- 303 Marsh GW, Stewart JS. Splenic function in adult coeliac disease. *Br J Haematol* 1970; 19: 445–57.
- 304 William BM, Thawani N, Sae-Tia S, *et al.* Hyposplenism: a comprehensive review. Part II: clinical manifestations, diagnosis, and management. *Hematology* 2007; 12: 89–98.
- 305 Schmitz F, Herzig KH, Stuber E, *et al.* On the pathogenesis and clinical course of mesenteric lymph node cavitation and hyposplenism in coeliac disease. *Int J Colorectal Dis* 2002; 17: 192–8.
- 306 Fanos V, Verlato G, Matti P, *et al.* Increased incidence of urinary tract infections in patients with coeliac disease. *Pediatr Nephrol* 2002; 17: 570–1.
- 307 Saalman R, Fallstrom SP. High incidence of urinary tract infection in patients with coeliac disease. *Arch Dis Child* 1996; 74: 170–1.
- 308 Ludvigsson JF, Olén O, Bell M, Montgomery SM. Coeliac disease and risk of sepsis. *Gut* 2008; 57: 1074–80.
- 309 Williams AJ, Asquith P, Stableforth DE. Susceptibility to tuberculosis in patients with coeliac disease. *Tubercle* 1988; 69: 267–74.
- 310 Ludvigsson JF, Wahlstrom J, Grunewald J, *et al.* Coeliac disease and risk of tuberculosis: a population based cohort study. *Thorax* 2007; 62: 23–8.
- 311 Shetty A, McKendrick M. TB and coeliac disease. *J Infect* 2004; 48: 109–11.
- 312 Holt PD, Tandy NP, Anstee DJ. Screening of blood donors for IgA deficiency: a study of the donor population of south-west England. *J Clin Pathol* 1977; 30: 1007–10.
- 313 Heneghan MA, Stevens FM, Cryan EM, *et al.* Celiac sprue and immunodeficiency states: a 25-year review. *J Clin Gastroenterol* 1997; 25: 421–5.
- 314 Cataldo F, Marino V, Ventura A, *et al.* Prevalence and clinical features of selective immunoglobulin A deficiency in coeliac disease: an Italian multicentre study. Italian Society of Paediatric Gastroenterology and Hepatology (SIGEP) and “Club del Tenue” Working Groups on Coeliac Disease. *Gut* 1998; 42: 362–5.
- 315 Not T, Faleschini E, Tommasini A, *et al.* Celiac disease in patients with sporadic and inherited cardiomyopathies and in their relatives. *Eur Heart J* 2003; 24: 1455–61.
- 316 Curione M, Barbato M, De Biase L, *et al.* Prevalence of coeliac disease in idiopathic dilated cardiomyopathy. *Lancet* 1999; 354: 222–3.
- 317 De Bem RS, Da Ro Sa Utiyama SR, Nishihara RM, *et al.* Celiac disease prevalence in Brazilian dilated cardiomyopathy patients. *Dig Dis Sci* 2006; 51: 1016–9.
- 318 Frustaci A, Cuoco L, Chimenti C, *et al.* Celiac disease associated with autoimmune myocarditis. *Circulation* 2002; 105: 2611–8.
- 319 Elfstrom P, Hamsten A, Montgomery SM, *et al.* Cardiomyopathy, pericarditis and myocarditis in a population-based cohort of inpatients with coeliac disease. *J Intern Med* 2007; 262: 545–54.
- 320 Ludvigsson JF, Welander A, Lassila R, *et al.* Risk of thromboembolism in 14,000 individuals with coeliac disease. *Br J Haematol* 2007; 139: 121–7.
- 321 Looker AC, Dallman PR, Carroll MD, *et al.* Prevalence of iron deficiency in the United States. *JAMA* 1997; 277: 973–6.
- 322 Corazza GR, Valentini RA, Andreani ML, *et al.* Subclinical coeliac disease is a frequent cause of iron-deficiency anaemia. *Scand J Gastroenterol* 1995; 30: 153–6.
- 323 Carroccio A, Iannitto E, Cavataio F, *et al.* Sideropenic anemia and celiac disease: one study, two points of view. *Dig Dis Sci* 1998; 43: 673–8.
- 324 Unsworth DJ, Lock RJ, Harvey RF. Improving the diagnosis of coeliac disease in anaemic women. *Br J Haematol* 2000; 111: 898–901.
- 325 Howard MR, Turnbull AJ, Morley P, *et al.* A prospective study of the prevalence of undiagnosed coeliac disease in laboratory defined iron and folate deficiency. *J Clin Pathol* 2002; 55: 754–7.
- 326 Ransford RA, Hayes M, Palmer M, *et al.* A controlled, prospective screening study of celiac disease presenting as iron deficiency anemia. *J Clin Gastroenterol* 2002; 35: 228–33.
- 327 Askling J, Linet M, Gridley G, *et al.* Cancer incidence in a population-based cohort of individuals hospitalized with celiac disease or dermatitis herpetiformis. *Gastroenterology* 2002; 123: 1428–35.
- 328 Cottone M, Termini A, Oliva L, *et al.* Mortality and causes of death in celiac disease in a Mediterranean area. *Dig Dis Sci* 1999; 44: 2538–41.
- 329 Card TR, West J, Holmes GK. Risk of malignancy in diagnosed coeliac disease: a 24-year prospective, population-based, cohort study. *Aliment Pharmacol Ther* 2004; 20: 769–75.
- 330 West J, Logan RF, Smith CJ, *et al.* Malignancy and mortality in people with coeliac disease: population based cohort study. *BMJ* 2004; 329: 716–9.
- 331 Collin P, Pukkala E, Reunala T. Malignancy and survival in dermatitis herpetiformis: a comparison with coeliac disease. *Gut* 1996; 38: 528–30.
- 332 Green PH, Fleischauer AT, Bhagat G, *et al.* Risk of malignancy in patients with celiac disease. *Am J Med* 2003; 115: 191–5.
- 333 Selby WS, Gallagher ND. Malignancy in a 19-year experience of adult celiac disease. *Dig Dis Sci* 1979; 24: 684–8.
- 334 Delco F, El-Serag HB, Sonnenberg A. Celiac sprue among US military veterans: associated disorders and clinical manifestations. *Dig Dis Sci* 1999; 44: 966–72.
- 335 Catassi C, Fabiani E, Corrao G, *et al.* Risk of non-Hodgkin lymphoma in celiac disease. *JAMA* 2002; 287: 1413–9.
- 336 Smedby KE, Akerman M, Hildebrand H, *et al.* Malignant lymphomas in coeliac disease: evidence of increased risks for lymphoma types other than enteropathy-type T cell lymphoma. *Gut* 2005; 54: 54–9.
- 337 Viljamaa M, Kaukinen K, Pukkala E, *et al.* Malignancies and mortality in

- patients with coeliac disease and dermatitis herpetiformis: 30-year population-based study. *Dig Liver Dis* 2006; **38**: 374–80.
- 338 Corrao G, Corazza GR, Bagnardi V, *et al.* Mortality in patients with coeliac disease and their relatives: a cohort study. *Lancet* 2001; **358**: 356–61.
- 339 Logan RF, Rifkind EA, Turner ID, *et al.* Mortality in celiac disease. *Gastroenterology* 1989; **97**: 265–71.
- 340 Green PHR, Stavropoulos SN, Panagi SG, *et al.* Characteristics of adult celiac disease in the USA: results of a national survey. *Am J Gastroenterol* 2001; **96**: 126–31.
- 341 Mustalahti K, Lohiniemi S, Kollin P, *et al.* Gluten-free diet and quality of life in patients with screen-detected celiac disease. *Eff Clin Pract* 2002; **5**: 105–13.
- 342 Hallert C, Granno C, Grant C, *et al.* Quality of life of adult coeliac patients treated for 10 years. *Scand J Gastroenterol* 1998; **33**: 933–8.
- 343 Hallert C, Lohiniemi S. Quality of life of celiac patients living on a gluten-free diet. *Nutrition* 1999; **15**: 795–7.
- 344 Sanchez-Albisua I, Wolf J, Neu A, *et al.* Coeliac disease in children with type 1 diabetes mellitus: the effect of the gluten-free diet. *Diabet Med* 2005; **22**: 1079–82.
- 345 Arato A, Korner A, Veres G, *et al.* Frequency of coeliac disease in Hungarian children with type 1 diabetes mellitus. *Eur J Pediatr* 2003; **162**: 1–5.
- 346 Westman E, Ambler GR, Royle M, *et al.* Children with coeliac disease and insulin dependent diabetes mellitus – growth, diabetes control and dietary intake. *J Pediatr Endocrinol Metab* 1999; **12**: 433–42.
- 347 Valletta E, Ulmi D, Mabboni I, *et al.* Early diagnosis and treatment of celiac disease in type 1 diabetes. A longitudinal, case-control study. *Pediatr Med Chir* 2007; **29**: 99–104.
- 348 Kaukinen K, Halme L, Collin P, *et al.* Celiac disease in patients with severe liver disease: gluten-free diet may reverse hepatic failure. *Gastroenterology* 2002; **122**: 881–8.
- 349 Tursi A, Giorgetti GM, Iani C, *et al.* Peripheral neurological disturbances, autonomic dysfunction, and antineuronal antibodies in adult celiac disease before and after a gluten-free diet. *Dig Dis Sci* 2006; **51**: 1869–74.
- 350 Chin RL, Latov N. Peripheral Neuropathy and Celiac Disease. *Curr Treat Options Neurol* 2005; **7**: 43–8.
- 351 Meloni G, Dore A, Fanciulli G, *et al.* Subclinical coeliac disease in schoolchildren from northern Sardinia. *Lancet* 1999; **353**: 37.
- 352 Mora S, Barera G, Beccio S, *et al.* Bone density and bone metabolism are normal after long-term gluten-free diet in young celiac patients. *Am J Gastroenterol* 1999; **94**: 398–403.
- 353 Corazza GR, Frisoni M, Vaira D, *et al.* Effect of gluten-free diet on splenic hypofunction of adult coeliac disease. *Gut* 1983; **24**: 228–30.
- 354 Trewby PN, Chipping PM, Palmer SJ, *et al.* Splenic atrophy in adult coeliac disease: is it reversible? *Gut* 1981; **22**: 628–32.
- 355 Curione M, Barbato M, Viola F, *et al.* Idiopathic dilated cardiomyopathy associated with coeliac disease: the effect of a gluten-free diet on cardiac performance. *Dig Liver Dis* 2002; **34**: 866–9.
- 356 Goel NK, McBane RD, Kamath PS. Cardiomyopathy associated with celiac disease. *Mayo Clin Proc* 2005; **80**: 674–6.
- 357 Amin R, Murphy N, Edge J, *et al.* A longitudinal study of the effects of a gluten-free diet on glycemic control and weight gain in subjects with type 1 diabetes and celiac disease. *Diabetes Care* 2002; **25**: 1117–22.
- 358 Hallert C, Sedvall G. Improvement in central monoamine metabolism in adult coeliac patients starting a gluten-free diet. *Psychol Med* 1983; **13**: 267–71.
- 359 Berg NO, Dahlqvist A, Lindberg T, *et al.* Correlation between morphological alterations and enzyme activities in the mucosa of the small intestine. *Scand J Gastroenterol* 1973; **8**: 703–12.
- 360 Yachha SK, Srivastava A, Mohindra S, *et al.* Effect of a gluten-free diet on growth and small-bowel histology in children with celiac disease in India. *J Gastroenterol Hepatol* 2007; **22**: 1300–5.
- 361 Molteni N, Bardella MT, Vezzoli G, *et al.* Intestinal calcium absorption as shown by stable strontium test in celiac disease before and after gluten-free diet. *Am J Gastroenterol* 1995; **90**: 2025–8.